

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**



JD

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁴ : C07C 47/11, 69/138, 87/08 C07C 87/29, 121/68, 147/14 C07C 149/273, C07D 207/263 C07D 231/12, 233/3, 233/42 C07D 233/60, 239/10, 239/22 C07D 239/36, 285/06, 471/02 A61K 31/38, 31/40, 31/415 A61K 31/505</p>	A1	<p>(11) International Publication Number: WO 87/06576</p> <p>(43) International Publication Date: 5 November 1987 (05.11.87)</p>
<p>(21) International Application Number: PCT/US86/00958</p> <p>(22) International Filing Date: 29 April 1986 (29.04.86)</p> <p>(71) Applicant (for all designated States except US): PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): SACCOMANO, Nicholas, A. [US/US]; 14 Woodland Terrace, Rural Route 4 Box 95, Westerly, Washington County, RI 02891 (US). VINICK, Fredric, J. [US/US]; 55 Trumbull Road, Waterford, New London County, CT 06385 (US).</p>	<p>(74) Agents: KNUTH, Charles, J. et al.; Pfizer Inc., Patent Department, 235 East 42nd Street, New York, NY 10017 (US).</p> <p>(81) Designated States: FI, HU, NO, SU, US.</p> <p>Published With international search report.</p>	
<p>(54) Title: CALCIUM INDEPENDENT cAMP PHOSPHODIESTERASE INHIBITOR ANTIDEPRESSANT</p> <div style="text-align: center; margin: 20px 0;">  <p>(I)</p> </div> <p>(57) Abstract</p> <p>Antidepressant agents having formula (I), wherein R¹ is a polycycloalkyl group; R² is methyl or ethyl, X is O or NH; and Y comprises a 5- or 6-membered heterocyclic ring having one or two nitrogens; or fused bicyclic heterocyclic rings having a total of three nitrogen atoms, one in each ring and one angular nitrogen.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				


-1-

CALCIUM INDEPENDENT CAMP PHOSPHODIESTERASE
INHIBITOR ANTIDEPRESSANT

Background of the Invention

Field of the Invention


This invention relates to antidepressant agents having the formula



wherein R^1 is a polycycloalkyl group; R^2 is methyl or ethyl and Y is a saturated or unsaturated 5- or 6-membered nitrogen-containing heterocyclic ring or a fused bicyclic heterocyclic ring having three nitrogen atoms. More particularly it relates to such compounds wherein Y is a saturated or unsaturated 5- or 6-membered heterocyclic ring having one or two nitrogen atoms; or fused bicyclic heterocyclic rings having one nitrogen atom in each ring and one angular nitrogen.

Description of the Prior Art

U.S. Patents 4,012,495 and 4,193,926, a continuation-in-part thereof, describe a series of 4-(polyalkoxyphenyl)-2-pyrrolidones having the formula (A)




-2-

wherein R_1 and R_2 are hydrocarbon radicals of up to 18 carbon atoms or optionally substituted (C_{1-5})alkyl groups; R_3 is H or OCH_3 ; R_4 is H, alkyl, aryl or acyl, and X is O or S, which compounds have neuropsychotropic activity. Examples of hydrocarbon R_1 and R_2 groups are, inter alia, cycloalkyl and cycloalkylalkyl, preferably of 3 to 7 carbon atoms. Compounds related to those of formula (A) but substituted at the 1-position of the pyrrolidone ring with a $-C(O)R$ group where R is alkyl, aryl, aralkyl, amino or substituted amino group are disclosed as neuropsychotropic agents in U.S. Patent 4,153,713.

A series of analogous compounds having formula (B)

15



(B)


20

wherein R_1 and R_2 can be as defined above in formula (A); R_3 is hydrogen, optionally substituted alkyl, alkenyl, aryl, aralkyl or acyl; R_4 and R_5 can be hydrogen; and X is O or S are described in British Patent 1,588,639. They are alleged to exhibit central depressive, antidopaminergic, antinociceptive and anticonvulsant actions and to have a certain similarity to neuroleptics.

30


-3-

U.S. Patent 4,308,278 discloses related compounds of formula (C) having anorexigenic activity



15 wherein R^1 is (C_{3-6}) cycloalkyl or benzyl; each of R^2 and R^3 is hydrogen or (C_{1-4}) alkyl; R^4 is R^2 or alkoxy carbonyl; and R^5 is hydrogen or alkoxy carbonyl.

U.S. Patent 3,636,039 and U.S. 3,923,833 a division thereof, disclose benzylimidazolidinones of formula (D) as hypertensive agents




30 wherein variables R_1 - R_4 are chosen from a variety of values including hydrogen and lower alkoxy.

-4-

Preparation of antihypertensive 1,4,5,6-tetrahydropyrimidines from appropriate benzaldehydes via intermediate glutaronitrile and glutaramide derivatives is described in U.S. 4,261,995.

Summary of the Invention

The antidepressants of this invention have the formula (I)



(I)

wherein R¹ is a polycycloalkyl group having from 7 to 11 carbon atoms;

R² is methyl or ethyl;

X is O or NH; and

Y comprises a 5- or 6-membered heterocyclic ring, especially a saturated or unsaturated 5- or 6-membered heterocyclic ring having one or two nitrogen atoms, said ring being optionally substituted with =O or =S, provided that when said optional group is present and the heterocyclic ring comprises one nitrogen atom the optional group is located on a carbon atom adjacent said nitrogen atom, and when the heterocyclic ring comprises two nitrogen atoms in a 1,3-position to one another said optional group is located on the carbon atom between the two nitrogen atoms; the nitrogen atoms of the heterocyclic rings which bear a hydrogen atom being optionally substituted with (C₁₋₅)alkyl, (C₂₋₅)alkenyl, (C₁₋₅)alkanoyl, benzyl, phenylethyl or benzoyl; 1,2,3-thiadiazolyl 2,2-dioxide, the nitrogen atoms of which are optionally substituted with (C₁₋₅)alkyl, (C₂₋₅)alkenyl, (C₁₋₅)alkanoyl, benzyl, phenylethyl or benzoyl; bicyclic heterocyclic moiety

-5-

rings containing a total of three nitrogen atoms, one in each of the rings and one common to each ring (angular nitrogen).

Also included in this invention are pharmaceutically acceptable acid addition salts of formula (I) compounds which have a basic nitrogen atom. Further, pharmaceutical compositions of formula (I) compounds and their use as antidepressant agents are embraced within the scope of this invention. Still further, processes for making formula (I) compounds and intermediates therefor, and the novel intermediates themselves are included in this invention. As will be recognized by those skilled in the art, stereocenters exist in several of the polycyclic (R^1) groups and heterocyclyl (Y) moieties of compounds of this invention. The racemic-diastereomeric mixtures and the individual optical isomers are also included in this invention.





As part of the present invention are certain compounds described herein which are especially useful as intermediates for preparation of formula (I) compounds. The compounds have formulae (IV), (V), (VII), (VIII), (IX), (XI), (XII), (XIII), (XIV) and (XVI) presented below. In the case of compounds of formulae (XI) and (XII), the tautomeric forms of said compounds are included in this invention, even though only one form, the ene form, is represented by said formulae.

Formula (I) compounds are characterized by a favorable therapeutic index and are significantly less emetic than are known compounds such as 4-[3-(cyclopentyloxy)-4-methoxyphenyl]-2-pyrrolidinone, which is described in U.S. Patent 4,193,926, and which is known by the non-proprietary name of rolipram.

-6-

Detailed Description of the Invention



More particularly this invention relates to formula (I) compounds wherein Y is




30

-7-


5



10



or



(o)

(p)

15

and R^1 , R^2 and X are as defined above for formula I compounds; R^3 is hydrogen, (C_{1-5}) alkyl, (C_{2-5}) alkenyl, benzyl or phenethyl; and R^4 is hydrogen, (C_{1-5}) alkyl, (C_{1-5}) alkanoyl, optionally having a basic group, or benzoyl.

Examples of R^1 as (C_{7-10}) polycycloalkyl are:

20

bicyclo[2.2.1]heptyl

bicyclo[2.2.2]octyl

bicyclo[3.2.1]octyl

tricyclo[3.2.1.0^{2,6}]decyltricyclo[3.3.1.1^{3,7}]decyl

indanyl .

Favored as values of R^1 are:

25

bicyclo[2.2.1]hept-2-yl

bicyclo[2.2.2]oct-2-yl

bicyclo[3.2.1]oct-3-yl

tricyclo[3.2.1.0^{2,6}]dec-4-yltricyclo[3.2.1.0^{2,6}]dec-8-yltricyclo[3.3.1.1^{3,7}]dec-2-yl

30

indan-2-yl.

Preferred values of R^1 are bicyclo[2.2.1]-hept-2-yl and indan-2-yl.

-8-

For each of the above categories the value of X as O and R² as methyl represent generally preferred compounds.

For a given value of R¹, favored values of Y are the 5-membered heterocyclyl rings, the bicyclic heterocyclic rings and the saturated 6-membered heterocyclic ring system. Preferred values of Y are the saturated 5- and 6-membered heterocyclic ring systems and the bicyclic heterocyclic ring systems (ring systems (a)-(d), (g), (l) and (m)-(p) above). Especially preferred are the heterocyclic ring systems (a), (g), (l) and (m)-(p).


From the standpoint of biological activity the exo isomer of a given compound of formula (I) is generally favored over the endo isomer because of greater potency. In actual practice in the case of formula (I) compounds having the asymmetric center in variables R¹ and/or Y, a mixture of the isomers of a given formula (I) compound is often favored because of relative ease of preparation of the mixture as compared to the pure isomers.

Products of this invention of formula (I) wherein X is O and Y represents heterocyclyl rings (a), (b) or (c) are prepared according to Sequence A:

25

30

-9-



In sequence A, conversion of (II) to (III) is accomplished by the Williamson synthesis involving nucleophilic substitution of a phenoxide ion for halide ion. The known methodology comprises transformation of a phenolic reactant (II) to a phenoxide ion by reaction

-10-

with a base, such as an alkali metal hydroxide or carbonate, in a reaction-inert solvent. By reaction-inert solvent is meant a solvent which does not react with starting materials, intermediates or product so as to adversely effect the yield of desired product. Suitable solvents are dimethylformamide, tetrahydrofuran, dioxan and bis(2-methoxyethyl)ether. The appropriate R^1 halide, preferably R^1Br or R^1Cl , is reacted with the phenoxide ion at an elevated temperature, e.g., up to about $150^\circ C.$, and the mixed ether (III) recovered by known procedures. In this reaction the use of the exo- or endo isomer of the polycycloalkyl halide results in production of an isomeric mixture of the corresponding ethers (III) which, if desired, can be separated by known procedures such as chromatography.

Alternatively, the mixed ethers of formula (III) are prepared by reacting the appropriate polycycloalkanol (R^1-OH) and phenol (II) in the presence of triphenylphosphine-diethyl azodicarboxylate as activator of the hydroxyl groups involved in the reaction. Reaction is conducted in a reaction-inert solvent such as tetrahydrofuran (THF) generally adding the diethyl azodicarboxylate at room temperature to a solution of polycycloalkanol, phenol and triphenylphosphine in THF. Upon completion of the addition, the reaction is refluxed until essentially complete and the mixed ether recovered by known procedures.

Still further, they are prepared from the appropriate polycycloalkanone by reacting it with catechol in a hydrocarbon solvent in the presence of p-toluenesulfonic acid under conditions which remove by-product water to produce the polycycloalkyl ketal. Lithium aluminum hydride/ $AlCl_3$ reduction of the ketal

-11-

affords the 2-polycycloalkyloxy phenol which is then brominated to give the 4-bromo-(2-polycycloalkyloxy)-phenol, which is then converted to the OR^2 ether at the phenolic group. Replacement of the bromo function by formyl is achieved by treating it with tert-butyl-lithium in THF followed by quenching with N,N-dimethyl-formamide. All of the above mentioned reactions are carried out according to known procedures. Other starting materials required for this invention, if not available, or not previously described in the art, are prepared by known procedures such as those described herein.

The aminonitriles or cyanoamines of formula (IV) are produced by the Strecker synthesis which involves replacement of the carbonyl oxygen of (III) with amino, or substituted amino, and cyano groups. The procedure comprises simultaneous reaction of aldehyde (III) with the appropriate amine (or ammonia) as its hydrochloride salt and sodium cyanide in a reaction-inert solvent such as ethanol. The reaction proceeds in good yield at ambient temperatures and the product (IV) is recovered by known methods.

Formula (V) compounds are obtained by reduction of formula (IV) compounds. As is well known reduction of nitriles to amines can be carried out by a variety of means including catalytic hydrogenation over a noble metal or Raney nickel catalyst, by hydrides such as lithium aluminum hydride, diisobutylaluminum hydride, sodium diethylaluminum hydride and sodium bis(2-methoxyethoxy)aluminum hydride. A favored reducing agent in this invention is diisobutylaluminum hydride, (known as DiBal-H). The reduction is conveniently carried out by adding a solution of the cyano amine (IV) in a reaction-inert solvent, e.g. toluene,

-12-


cyclohexane, diethyl ether, tetrahydrofuran, heptane, hexane and dichloromethane, to a solution of DiBal-H in a reaction-inert solvent, preferably in the same solvent as used for the cyanoamine. The reaction is conducted at a low temperature, e.g., below -50°C. during addition of the cyanoamine to the reducing agent, and for a period of time (2-4 hours, following completion of addition. It is then warmed slowly to about 0°C., excess reducing agent destroyed, and the product recovered by standard methods.

Cyclization of (V) to (I-a) wherein Y is a 2-imidazolidinon-5-yl group is achieved by treating the diamine (V) in a reaction-inert solvent with N,N'-carbonyldiimidazole or 1,1-carbonyldi-1,2,4-triazole in tetrahydrofuran or tetrahydrofuran-benzene at from 20° to 65°C. Phosgene can be used but because of its toxicity relative to the aforementioned cyclizing reagents, and the need for higher temperatures for cyclization is not favored. The corresponding thione of formula (I-b) wherein Y is a 2-imidazolidinthion-5-yl group is prepared in like manner but using N,N'-thiocarbonyldiimidazole as cyclizing agent. The formula (I-c) compounds wherein Y is a 1,2,3-thiadiazol-5-yl 2,2-dioxide are prepared by using sulfamide as cyclizing agent. Reaction is conducted in pyridine at reflux temperature until complete and the product recovered by known methods.

30

-13-

Compounds wherein Y of formula (I) is heterocyclyl ring (f) and X is O are prepared according to sequence B:




-14-

The formula (VI) reactants are prepared from compounds of formula (III) by procedures known to those skilled in the art. A convenient procedure comprises oxidation of (III) by means of Jones reagent (chromic acid and sulfuric acid in water). The oxidant is added to the benzaldehyde reactant (III) in acetone at ambient temperature and the acid (VI) recovered by standard methods. The acid (VI) is next esterified by known procedures to, desirably, a lower alkyl ester and preferably the methyl ester. The ester is then reacted with about two equivalents of sodium methylsulfinylmethide (formed by reaction of sodium hydride with dimethyl sulfoxide) in tetrahydrofuran at from about 0°-10°C. to provide the beta-ketosulfoxide (VII). Isomerization of ketosulfoxide (VII) with aqueous acid results in rearrangement (Pummerer) of the ketosulfoxide to the hemimercaptal (VIII). Treatment of the hemimercaptal with cupric acetate monohydrate in a reaction-inert solvent such as chloroform affords the alpha-keto aldehyde (IX). The keto aldehydes of formula (IX) are readily transformed to formulae (I-f) compounds by reaction with formaldehyde and ammonium hydroxide in a reaction-inert solvent such as a lower alkanol, especially ethanol, at from about 20°C. to 50°C. to provide the imidazoles. They are conveniently isolated as their acid addition salts.


Formula (I) compounds wherein X is O and Y is heterocyclyl rings (e), (g), (h), (i), (m) or (o) above are prepared according to sequence C:

30

-15-



-16-



In sequence C reactants having formula (X) are prepared from formula (III) compounds by known procedures comprising reduction of the aldehyde function to hydroxymethyl by, for example, sodium borohydride, conversion of the alcohol to the corresponding bromo derivative by means of triphenylphosphine-carbon tetrabromide, and reaction of the thus-produced bromo derivative with alkali metal cyanide to afford (X). Reaction of (X) with ethyl formate in an aprotic reaction-inert solvent such as an aromatic or aliphatic hydrocarbon such as benzene, toluene or hexane; an ether such as dioxane, tetrahydrofuran, bis(2-methoxyethyl)ether; and in the presence of a base of sufficient strength, such as sodium hydride, to form an anion at the active methylene group provides formula (XI) compounds.

Conversion of (XI) to (XII) is accomplished by hydrogenation over Raney nickel in a reaction-inert solvent such as a lower alkanol, at pressures of from about 15 to about 100 psi (about 1 to about 7 kg/sq cm) at ambient temperature. Other catalysts such as noble metal catalysts can be used. However, their use may

-17-

bring about reduction of the formyl group along with the nitrile group and is avoided. Cyclization of (XII) to (I-e) is readily carried out by treating the enamino-aldehyde (XII) with hydrazine in a lower alkanol solvent at about 50° to 100°C. The pyrazole derivative (e) is isolated by procedures known to the skilled artisan.

The 6-membered heterocyclyl structure (I-g) is produced by reaction of the enamino-aldehyde (XII) with urea in the presence of hydrochloric acid, or other strong acid, and in a reaction-inert solvent such as a lower alkanol or ethers such as dioxane, tetrafuran, bis(2-methoxyethyl)ether and 1,2-dimethoxyethane. The pyrimidinone (I-g) is then reduced to obtain (I-h) and/or (I-i). Reduction of (I-g) with Raney nickel in a reaction-inert solvent such as those enumerated above at a temperature of from about 50°C. to refluxing temperature of the chosen solvent, affords the tetrahydropyrimidinone (I-h). Further reduction over Raney nickel provides the hexahydropyrimidinone (I-i). In each of the above reductions a hydrogen pressure of from about 15 to about 100 psi is productive of satisfactory yields. Other catalysts such as noble metal catalysts can, of course, be used for these reductions, if desired.

Alternatively, compounds of formula (I-i) are prepared by reacting the appropriate 3-R¹O-4-R²O-benzaldehyde of formula (III) with cyanoacetic acid in a reaction-inert solvent, e.g. pyridine, in the presence of piperidine, morpholine, piperazine, or other base which is more basic than pyridine, at from ambient temperature to about 120°C.; and preferably from 50° to 100°C. The 3-(3-R¹O-4-R²O phenyl)glutaro-

-18-


nitrite is isolated by known procedures as, for example, by pouring the reaction mixture in water and extracting the product with a suitable solvent, such as ethyl acetate. The glutaronitrite derivative is then converted to the corresponding glutaramide derivative by treatment in a reaction-inert solvent, e.g. aqueous acetone, with hydrogen peroxide and sodium carbonate at 0°C. The reaction mixture is slowly warmed to ambient temperature and stirred until complete. The diamide is recovered by concentration and extraction of the reaction mixture. The 3-(3-R¹O-4-R²O)glutaramide is then cyclized by reaction in a reaction-inert solvent, e.g. pyridine, using lead tetraacetate at ambient temperature. The product is recovered by extraction.

Compounds of formulae (I-m) and (I-o) are produced from the enaminoaldehyde (XII) by reacting it with 3-aminopyrazole or a 2-aminoimidazole, respectively in the presence of a mineral acid. Ethanol is favored as a solvent although reaction-inert solvents such as are enumerated above for preparation of (I-g) can be used. The reaction is conducted at temperatures of from about 50°C. to the reflux temperature of the chosen solvent.

Formula (I) compounds wherein the heterocyclyl ring Y is (d) above are prepared as outlined in sequence D:

30

-19-



20

Knoevenagel condensation of aldehyde (III) with a di(C_{1-5})alkyl malonate, e.g., diethylmalonate, in a reaction-inert solvent, e.g., benzene, toluene, xylene, hexane, cyclic ethers, bis(2-methoxyethyl)ether, 1,2-dimethoxy ethane, in the presence of piperidine at 50°-150°C. yields the unsaturated diester (XIII).

25

Reaction of the diester in ethanol or other reaction-inert solvent with sodium cyanide at ambient temperature gives the cyano ester (XIV). Reduction of the cyano ester in acetic acid using a noble metal catalyst, e.g., platinum oxide, affords the

30

corresponding amino acid ester which is then cyclized to the corresponding alpha-carbethoxy lactam by heating in an aromatic hydrocarbon or ether solvent such as those mentioned above for the Knoevenagel condensation.

-20-

Saponification of the carbethoxy lactone with
ethanolic-alkali metal hydroxide at reflux and
neutralization of the sodium salt thus produced
provides the alpha-carboxy lactam which is thermally
5 decarboxylated at about 180°C. to afford the
pyrrolidone (I-d).

Sequence E below outlines preparation of formula
(I) compounds wherein X is O and Y is heterocyclyl
rings (j), (k), (l), (n) and (p):

10


15

20

25

30


-21-



-22-

In this sequence a formula (III) compound is converted to carbinol (XV) by the well known Grignard reaction. The carbinol is then oxidized by means of Jones' reagent under conditions described above for Sequence B, formula (VI) compounds, to afford the corresponding ketone. Treatment of the ketone in tris-dimethylaminomethane, a reagent for introducing the dimethylaminoethylidene group, at the reflux temperature provides compounds of formula (XVI). This versatile reactant serves as the key building block for heterocyclyl compounds (I-j), (I-k), (I-l), (I-n), and (I-p) of this invention.

Reaction of (XVI) with urea in ethanolic hydrochloric acid affords (I-j) or a related compound wherein the heterocyclyl moiety is 4-(4-hydroxy-1,2,3,4-tetrahydro-2-pyrimidinone) of the formula



(ja)

or a mixture of said two types of compounds. The product of the cyclization of urea with (XVI) depends upon the reaction time. Conduct of the reaction at or near reflux for about one hour affords principally the hydrated product wherein the heterocyclyl moiety is (ja). A heating period of about two hours yields a mixture of the two products in roughly equal amounts. A longer reaction period, e.g., 4 to 6 hours, gives (I-j) compounds, dehydrated derivatives of the (I-ja) compounds as major product.

-23-

Reduction of (I-j) compounds, e.g., with Raney nickel or a noble metal catalyst affords compounds of formula (I-k) or (I-l). The particular product depends upon the reduction conditions as those skilled in the art will recognize.


Compounds of formula (I-j) are produced from the precursor hydrated compounds wherein Y is (ja) by heating in ethanolic hydrochloric acid until removal of the elements of water is complete.

Compounds of formula (I-p) and (I-n) are readily obtained by reacting formula (XVI) compounds with 2-aminoimidazole or 3-aminopyrazole, respectively, under conditions corresponding to those presented above for preparation of (I-o) and (I-m).

Compounds of formula (I) wherein XR^1 is $-\text{NH}-\text{R}^1$ are prepared by procedures similar to those presented in the above reaction sequences. The difference resides in use of the appropriate 3-nitro-4- OR^2 benzaldehyde, 3-nitro-4- OR^2 benzylcyanide or 3-nitro-4- OR^2 -benzoic acid, as starting material in place of the 3- OR^1 -4- OR^2 diether reactants of sequences A-E. Said reactants are put through the above reaction sequences to provide compounds wherein the $-\text{XR}^1$ group of formula (I) compounds is replaced by nitro. Reduction of said nitro group to amino by known procedures, as by use of platinum oxide/hydrogen affords the corresponding amino derivative. Reductive alkylation of the amino derivative with the appropriate polycycloalkyl ketone according to procedures known to those skilled in the art affords the formula (I) compound in which the amino group NH_2 has been converted to $-\text{NHR}^1$.

-24-

Also of interest for the same utilities and in the same manner are formula I compounds wherein Y is one of the following heterocyclyl moieties:



Z is hydrogen, C₁₋₄ alkoxy, Cl; and

15 n is 1 or 2. Said compounds are prepared according to the general procedure described in Indian J. Chem., 18B, 428 (1979). The procedure comprises reacting equimolar amounts of the 3-R¹O-4-R²O benzaldehyde of formula (III) above and the appropriate
20 ortho-diamine of the 6-membered ring component of the fused ring system (e.g., 1,2-diaminobenzene) at 100°C. in nitrobenzene. The nitrobenzene serves as both solvent and oxidizing agent. Reaction is generally complete in 2-3 hours. The product is recovered by
25 known methods, including flash column chromatography over silica gel.

The acid addition salts of formula (I) compounds having a basic group are readily prepared by adding a stoichiometric amount of the appropriate acid to a
30 formula (I) compound in a solvent, preferably one in which the formula (I) compound is at least partially soluble. The acid addition salt, if soluble in the solvent system is recovered by evaporation of the solvent or by addition of a non-solvent for the salt to precipitate it from the reaction solvent.

-25-

The compounds of this invention having formula (I) function as calcium independent c-AMP phosphodiesterase inhibitors and are useful as antidepressants. Their activity as calcium independent c-AMP phosphodiesterase inhibitors is determined by the method of Davis, Biochimica et Biophysica Acta. 797, 354-362 (1984). In this procedure, calcium-independent and dependent phosphodiesterases (IPDE and DPDE, respectively) are prepared from cerebral cortices of female Sprague-Dawley rats by first homogenizing the brain tissue in a pH 7.5 20 mM Tris-HCl buffer also containing 1 mM $MgCl_2$, 3 mM 2-mercaptoethanol, and 0.1 mM EGTA (ethyleneglycol-bis-(beta-aminoethyl ether)-N,N'-tetraacetic acid). The homogenate is centrifuged at 105,000 x g for 60 minutes, and the supernatant fluid containing the enzymes is passed through a column of Sephadex G-200 to separate IPDE from DPDE. The two phosphodiesterases are each further purified by affinity chromatography on a column of calmodulin-Sepharose.

Phosphodiesterase activity is determined using 0.1 ml reaction mixture containing Tris-HCl pH 7.5 buffer (5 μ mol), $MgCl_2$ (0.5 μ mol), and [3H]cAMP (New England Nuclear, NET-275). The final concentration of cAMP is 1.0 μ M (containing 400,000 dpm of [3H]cAMP). Ten μ l of vehicle or inhibitor solution and 10 μ l of fresh IPDE or DPDE or the respective boiled enzymes are added to 80 μ l of substrate in the Tris-HCl/ $MgCl_2$ buffer. The reaction mixtures are incubated for 8 minutes at 37°C. and placed in a hot water bath for 2 minutes to stop hydrolysis of cAMP. Carrier 5'-AMP (0.5 ml of 0.5 mM 5'-AMP in 0.1 M Hepes (N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid)-0.1 M NaCl pH 8.5 buffer) are added, and the contents of the incubation tubes placed

-26-

on columns of polyacrylamide-boronate affinity gel (BIO-RAD Affi-Gel 601 Boronate Gel). The unreacted [3H]cAMP eluted from the gel with 7.5 ml of the 0.1M Hepes-NaCl buffer. The [3H]cAMP eluted from the gel with 7.5 ml of the 0.1M Hepes-NaCl buffer. The [3H]5'-AMP product is eluted with 7 ml of 50 mM Na acetate buffer pH 4.8. One-ml aliquots of the latter eluates are counted in a liquid scintillation counter to determine their content of radioactive 5'-AMP.

When used for the treatment of depression and other various neurological and psychic disorders, and characterized by withdrawal, anxiety, thought-disturbances and delusion, they are used as is or in the form of pharmaceutical compositions comprising a formula (I) compound and pharmaceutically-acceptable carriers or diluents. For oral administration, the preferred route for administering said compounds, suitable pharmaceutical carriers include inert diluents or fillers, thereby forming dosage forms such as tablets, powders, capsules, and the like. These pharmaceutical compositions can, if desired, contain additional ingredients such as flavorings, binders, excipients and the like. For example, tablets containing various excipients, such as sodium citrate, are employed, together with various disintegrants such as starch, alginic acid and certain complex silicates, together with binding agents such as polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often useful for tabletting purposes. Solid compositions of a similar type may also be employed as fillers in soft and hard filled gelatin capsules. Preferred materials therefor include

-27-

lactose or milk sugar and high molecular weight polyethylene glycols.

For oral administration, the daily dose of active agent of formula (I) is from about 0.1 mg to about 10 mg., and for parenteral administration, preferably i.v. or i.m., from about 0.01 mg. to about 5 mg. The prescribing physician, of course, will ultimately determine the appropriate dose for a given human subject dependent upon factors such as the severity of the patient's symptoms and the patient's response to the particular drug.

The antidepressant activity of the compounds of this invention is determined by the behavioral despair paradigm described by Porsolt et al., Arch. Int. Pharmacodyn. 227, 327-336 (1977).

In this procedure a depressed state is induced in mice by forcing them to swim in a narrow water-containing cylinder from which there is no escape. The procedure involves injecting a mouse per os with the test compound and then (30 minutes post-injection) placing it in a standard 1 liter glass beaker containing 800 ml of 25 degree Centigrade water.

An observer then rates the animal's mobility (0 = mobile; 1 = immobile) every 30 seconds for 5 minutes beginning 2 minutes after being placed in the water. Male CD (Charles River) mice (10 per treatment) weighing 20-25 g are used as subjects. The compounds are administered in a solution (vehicle) containing 0.9% saline (90%), dimethyl sulfoxide (5%), and emul 4 (5%). All drugs are injected in a volume of 10 ml/kg. A vehicle treated mouse typically has a swim score of 9, while an antidepressant drug reduces the magnitude of immobility, resulting in a decrease in the swim score.

-28-

The second procedure is the method of Koe et al., J. Pharmacol. Exp. Therap. 226, 686-700 (1983) which comprises determination of the ability of a test drug to counteract reserpine hypothermia in mice. In this procedure, mice are placed in a room with an ambient temperature of 20°C. The mice are individually housed in plastic chambers with a cardboard bottom, injected with reserpine (1.0 mg/kg s.c.), and retained at 18-19°C. for 18 hours. Their rectal temperatures are then ascertained, immediately after which they receive saline or drug treatment. Rectal temperatures are again measured, usually at 1, 2 and 4 hours after this second injection. Results are presented as the mean increase in reserpine-depressed temperature, expressed either as a percentage or an absolute increase. Typically, reserpine-pretreated mice given vehicle exhibit rectal temperatures averaging about 20-22°C. 4 hours after vehicles. Treatment with desipramine (10 mg/kg p.o.), a known antidepressant, yields temperatures averaging about 30-33°C. (about a 40-50% increase). Administration of formula (I) compounds brings about an increase in rectal temperature of the test mice.

Formula (I) compounds are less emetic than is rolipram and afford, therefore, an advantage over rolipram. Their emetic behavior is determined by administering to dogs the test drug dissolved in ethanol at 10 mg/ml and dose diluted to final volume chosen with distilled water, the ethanol content of the final solution not exceeding 10%. The drug solution is administered via oral gavage in a constant volume of 2 ml/kg body weight and the dogs then observed for emesis. If emesis occurs, the latency (the time from injection time to emesis) is recorded. If no emesis

-29-

occurs within 30 minutes following administration of drug, a higher dose is administered to the next subject. A starting dose of 100 ug/kg was used based on the minimal effective dose of rolipram.

5 The following examples and preparations are provided solely for further illustration. The following abbreviations for peak shapes are used in reporting ¹H-nmr values: bs, broad singlet; s, singlet; d, doublet; t, triplet; q, quartet, m, multiplet. In the Examples and Preparations no effort
10 was made to optimize the yield of any given reaction.

15

20

25

30

-30-

EXAMPLE 1alpha-N-Methylamino-3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxybenzene Acetonitrile

The bicyclo[2.2.1]hept-2-yl ethers of Preparation 2 (3.5 g, .014 mol) are dissolved in 50 ml of ethanol and to it is added sodium cyanide (.736 g, .015 mol) and methylamine hydrochloride (1.0 g, .015 mol) and this is stirred for 18 hours at room temperature. The reaction mixture is diluted with saturated sodium bicarbonate solution and is extracted with, ethyl ether (3 x 30 ml). The combined organic layers are washed with saturated sodium chloride solution (3 x 30 ml), dried over anhydrous sodium sulfate, filtered and concentrated in vacuo to yield 4.15 g (~100%) of the cyanoamine as a clear oil. The cyanoamines are obtained as a 7:3 endo:exo mixture of bicycloalkyl ethers.

^1H Nmr (300 MHz, CDCl_3): delta 7.2-6.8 (m, 3H), 4.72 (bs, 1H), 4.6-4.7 (m, .7H) [endo], 4.2-4.3 (m, .3H) [exo], 3.9 (bs, 3H), 3.5 (m, 1H), 2.6 (bs, 3H), 2.7-1.0 (m, 10H).

In like manner the following conversions are carried out:

3-(Endo-tricyclo[3.2.1.0^{2,6}]dec-8-yloxy)-4-methoxybenzaldehyde is converted to the corresponding aminonitrile in 89.4% yield;

^1H NMR (300 MHz, CDCl_3): delta 7.1-6.81 (m, 3H), 4.7 (bs, 1H), 4.6 (m, 1H), 3.9 (s, 3H), 2.6 (s, 3H), 2.8-1.0 (m, 14H);

30

-31-

3-(exo-tricyclo[3.2.1.0^{2,6}]dec-4-yloxy)-4-methoxybenzaldehyde in 97.2% yield to the corresponding aminonitrile.

5 ¹H NMR (300 MHz, CDCl₃): delta 7.1 (m, 2H), 6.9 (m, 1H), 4.9 (m, 1H), 4.75 (s, 1H), 3.9 (s, 3H), 2.65 (s, 3H), 2.6 (m, 2H), 2.3-1.2 (m, 12H);

3-exo-benzobicyclo[2.2.1]hept-2-yloxy)-4-methoxybenzaldehyde in approximately quantitative yield to the corresponding aminonitrile:

10 ¹H-NMR (300 MHz, CDCl₃): delta 7.4-6.9 (m, 7H), 4.6 (bs, 1H), 4.4 (m, 1H), 3.9 (s, 3H), 3.6 (bs, 1H), 3.4 (bs, 1H), 3.35 (bs, 1H), 2.5 (bs, 3H), 2.25 (m, 1H), 1.95 (m, 3H);

15 3-(endo-benzobicyclo[2.2.1]hept-2-yloxy)-4-methoxybenzaldehyde in quantitative yield to the corresponding aminonitrile;

20 ¹H NMR (300 MHz, CDCl₃): delta 7.2-6.95 (m, 6H), 6.8 (m, 1H), 5.1 (m, 1H), 4.7 (bs, 1H), 3.7 (m, 1H), 3.61 (bs, 3H), 3.2 (m, 1H), 2.6 (s, 3H), 2.5 (m, 1H), 1.9 (m, 1H), 1.8 (m, 1H), 1.25 (m, 1H);

3-endo-tricyclo[3.2.1.0^{2,6}]dec-8-yloxy)-4-methoxybenzaldehyde to the corresponding aminonitrile in 95.4% yield:

25 ¹H NMR (300 MHz, CDCl₃): delta 7.0-6.7 (m, 3H), 4.7 (m, 1H), 4.65 (bs, 1H), 3.87, 3.85 (s, 3H), [2-methoxyls], 2.5 (bs, 3H), 2.2-2.5 (m, 14H).

-32-

EXAMPLE 22-Methylamino-2-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]ethylamine

To a 1 liter flame dried round bottom flask is
5 added di-isobutyl aluminum hydride (DiBal-H, 46.6 ml of
a 1.5 molar solution in toluene, .07 mol) and 150 ml of
dry toluene. The hydride solution is cooled to -78°C.
and to it is added dropwise the cyanoamine of Example 2
(4.00 g, .014 mol) as a solution in 250 ml of dry
10 toluene over a 1 hour period. The reaction is stirred
at -78°C. for 2 hours and is warmed slowly to 0°C where
it is quenched slowly by the dropwise addition of a
saturated solution of sodium potassium tartrate (10 ml).
When gas evolution is no longer evident an additional 40
15 ml of the tartrate solution is carefully added and the
reaction is warmed to room temperature. The reaction
slurry is diluted with 150 ml ethyl ether and the
aqueous layer is extracted with ethyl ether (2 x 50 ml).
The combined ethereal layer is washed with saturated
20 tartrate solution (2 x 50 ml), water (2 x 50 ml) and
saturated sodium chloride solution (2 x 50 ml). The
organics are dried over anhydrous sodium sulfate,
filtered and concentrated in vacuo to yield 3.19 g
(78.5%) of the diamine as a light yellow oil. The
25 diamines are obtained as a 7:3 endo to exo mixture of
bicyclo[2.2.1]hept-2-yloxy ethers.

30

-33-

In like manner, the following 2-methylamino-2-[3-(R¹O)-4-methoxyphenyl]ethylamines are prepared from appropriate reactants:

R¹O = endo-benzobicyclo[2.2.1]hept-2-yloxy in 39%

yield:

¹H NMR (300 MHz, CDCl₃): delta 7.2-6.7 (m, 7H), 5.2 (m, 1H), 3.7 (m, 1H), 3.6 (s, 3H), 3.2 (m, 2H), 2.85 (m, 2H), 2.5 (m, 1H), 2.4 (s, 3H), 1.9 (m, 1H), 1.8 (m, 1H), 1.2 (m, 1H);

R¹O = exo-benzobicyclo[2.2.1]hept-2-yloxy in 94.5%

yield;

R¹O = endo-tricyclo[3.2.1.0^{2,6}]dec-4-yloxy in 96%

yield:

¹H NMR (300 MHz, CDCl₃): delta 6.8-6.7 (m, 3H), 4.75 (m, 1H), 3.78 (s, 1H), 3.38 (m, 1H), 2.8 (m, 2H), 2.3 (s, 3H), 2.4-.9 (m, 14H).

¹H NMR (300 MHz, CDCl₃): delta 6.7-7.0 (m, 3H), 4.6 (m, .7H), 4.2 (m, .3H), 3.8 (bs, 3H), 3.64 (bs, 1H), 3.33 (m, 1H), 2.7-2.85 (m, 2H), 2.3 (bs, 3H), 2.5-1.0 (m, 10H).

In like manner the following diamines are prepared from appropriate reactants: 2-methylamino-2-[3-(endo-tricyclo[3.2.1.0^{2,6}]dec-8-yloxy)-4-methoxyphenyl]ethylamine in approximately quantitative yield;

¹H NMR (300 MHz, CDCl₃): delta 7.0-6.8 (m, 3H), 5.7 (m, 1H), 3.9 (s, 3H), 3.5 (m, 1H), 3.0-1.0 (m, 19H).

2-Methylamino-2-[3-(endo-tricyclo[3.2.1.0^{2,6}]dec-4-yloxy)-4-methoxyphenyl]ethylamine in 76.62% yield.

-34-

EXAMPLE 31-Methyl-5-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]-2-Imidazolidinone

5 The diamines of Example 2 (3.19 g, .011 mol) are dissolved in dry tetrahydrofuran and treated with 1,1-carbonyldiimidazole (2.67 g, .0165 mol). The reaction is stirred for 24 hours at room temperature. The reaction mixture is poured into 50 ml of water and extracted with ethyl acetate (2 x 30 ml). The combined
10 organics are washed with 1N sodium hydroxide solution (2 x 30 ml), 1N hydrochloric acid (2 x 30 ml), water (2 x 30 ml), and saturated salt solution (3 x 30 ml). The organics are dried over anhydrous sodium sulfate, filtered and concentrated in vacuo to a clear oil which
15 is triturated with ether to yield 622 mg (17.8%) of the imidazolidinone as a white solid. MP: 142-144°C. The material represents a mixture of 2 pairs of diastereoisomers.

¹H NMR (250 MHz, CDCl₃): delta 6.7-6.85 (m, 3H),
20 5.7 (bs, 1H), 4.6 (m, .7H), 4.4 (m, 1H), 4.2 (m, .3H), 3.82 (s, 2.1H), 3.80 (s, .9H), 3.68 (dd, J=11.5Hz, J=8Hz), 3.21 (dd, J=11.5, J=8.1Hz), 2.6 (s, 3H), 2.6-1.0 (m, 10Hz).

25

30

-35-

^{13}C NMR (63 MHz , CDCl_3): delta 163.2, 149.8, 148.7, 131.7, 119.3, 113.2, 112.3, 112.2, 112.0, 78.9, 78.8, 62.84, 62.81, 56.15, 56.0, 47.5, 41.1, 40.5, 39.9, 37.2, 37.1, 36.7, 35.4, 35.3, 29.4, 28.74, 28.70, 28.3, 24.2, 20.7 (30 lines).

MS: M^+ = 316.0, 222.1, 95

Similarly, the following imidazolidinones are prepared by cyclization of appropriate diamines:
1-methyl-5-[3-(endo-tricyclo[3.2.1.0^{2,6}]dec-8-yloxy)-4-methoxyphenyl]-2-imidazolidinone in 20.7% yield.

MP = 149-152°C.

^1H NMR (300 MHz , CDCl_3): delta 6.9-6.8 (m, 3H), 5.15 (bs, 1H), 4.65 (m, 1H), 4.5 (m, 1H), 3.9 (s, 3H), 3.75 (m, 1H), 3.25 (m, 1H), 2.7 (s, 3H), 2.8-1.0 (m, 14H).

HRMS 356.2120 Calcd. for $\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_3$ 356.2099.

Analysis:

	Calcd	Act.
C	70.76	70.72
H	7.92	7.86
N	7.86	7.79

1-methyl-5-[3-(exo-tricyclo[3.2.1.0^{2,6}]dec-4-yloxy)-4-methoxyphenyl]-2-imidazolidinone in 11.42% yield.

^1H NMR (300 MHz , CDCl_3): delta 6.8 (m, 3H), 6.1 (bs, 1H), 4.8 (m, 1H), 4.4 (m, 1H), 3.8 (bs, 3H), 3.7 (m, 1H), 3.2 (m, 1H), 2.6 (s, 3H), 3.6-1.2 (m, 14H).

^{13}C NMR (75.43 MHz , CDCl_3): delta 163.3, 150.1, 148.2, 131.9, 119.6, 113.6, 112.1, 83.4, 56.0, 47.5, 43.2, 42.3, 40.6, 32.7, 29.6, 28.7, 23.2 (17 lines).

Analysis Calcd. for $\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_3$:

C, 70.75; H, 7.91; N, 7.86.

C, 68.76; H, 7.61; N, 8.35.

-36-

1-methyl-5-[3-(endo-benzobicyclo[2.2.1]hept-2-yloxy)-4-methoxyphenyl]-2-imidazolidinone in 13.74% yield:

5 ^1H NMR (300 MHz, CDCl_3): δ 7.2-7.05 (m, 4H), 6.8 (m, 3H), 5.1 (m, 1H), 4.85 (bs, 1H), 4.45 (m, 1H), 3.7 (m, 1H), 3.68 (m, 1H), 3.4 (m, 1H), 3.25 (m, 1H), 2.65 (s, 1H), 2.45 (m, 1H), 1.9 (m, 1H), 1.8 (m, 1H), 1.25 (m, 1H);

10 1-methyl-5-[3-(exo-benzobicyclo[2.2.1]hept-2-yloxy)-4-methoxyphenyl]-2-imidazolidinone in about 17% yield:

15 ^1H NMR (300 MHz, CDCl_3): δ 7.2-7.1 (m, 4H), 7.0-6.9 (m, 3H), 4.5 (m, 1H), 4.4 (m, 1H), 3.9 (s, 1H), 3.75 (m, 1H), 3.6 (bd, 1H), 3.4 (bs, 1H), 3.25 (m, 1H), 2.67 + 2.65 (s, 3H), 2.25 (m, 1H), 1.95 (m, 3H).

Analysis Calcd. for $\text{C}_{22}\text{H}_{24}\text{N}_2\text{O}_3$:

C, 72.50; H, 6.63; N, 7.68.

Found: C, 71.73; H, 6.78; N, 7.28.

20 1-methyl-5-[3-(endo-tricyclo[3.2.1.0^{2,6}]dec-4-yloxy)-4-methoxyphenyl]-2-imidazolidone in 7.2% yield:

^1H NMR (300 MHz, CDCl_3): δ 6.8-6.7 (m, 3H), 5.75 (bs, 1H), 4.75 (bs, 1H), 4.42 (m, 1H), 3.83 (s, 3H), 3.72 (m, 1H), 3.28 (m, 1H), 2.65 (s, 3H), 2.3-.9 (m, 14H).

25 ^{13}C NMR (75.4 MHz, CDCl_3): δ 163.2, 150.8, 147.8, 131.8, 120.0, 114.5, 112.4, 82.5, 62.8, 56.2, 47.6, 46.2, 40.4, 37.7, 31.8, 28.8, 28.5 (17 lines).

Analysis Calcd. for $\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_3$:

C, 70.75; H, 7.91; N, 7.86.

30 Found: C, 69.74; H, 7.93; N, 7.48.

-37-

EXAMPLE 4

1-n-Butyl-3-Methyl-4-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]-2-Imidazolidinone

5 The imidazolidinone of Example 3 (.5 g, 1.58 mmol) is dissolved in 20 ml, THF and 5 ml DMG, treated with sodium hydride (41 mg, 1.73 mmol), cooled to 0°C. and treated with n-iodobutane (.581 g, 3.15 mmol). The reaction is warmed slowly to room temperature and stirred for 24 hours. It is then diluted with water, and extracted with ether. The organic layer is washed
10 with water, brine, dried over $MgSO_4$, filtered and concentrated in vacuo. The product is triturated with ether to afford 211 mg (35.8%) of the product. This material is 7:3 endo/exo isomer mixture.

15 Analysis Calcd. for $C_{22}H_{32}N_2O_3$:
C, 70.93; H, 8.66; N, 7.52.
Found: C, 69.95; H, 8.66; N, 7.43.

EXAMPLE 5

20 1-Acetyl-3-Methyl-4-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]-2-Imidazolidinone

Repetition of the procedure of Example 4, but using an appropriate amount of acetyl chloride in place of n-iodobutane affords the title compound in 29.3% yield.

Analysis (as hemihydrate):

25 Calcd. for $C_{20}H_{26}N_2O_4 \cdot \frac{1}{2}H_2O$:
C, 65.44; H, 7.41; N, 7.63.
Found: C, 65.36; H, 7.29; N, 7.00.

30

-38-

EXAMPLE 61,3-Dimethyl-4-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]-2-Imidazolidinone

Repetition of the procedure of Example 4 but using a stoichiometric amount of iodomethane in place of n-iodobutane provides the title compound in 56.3% yield.

Analysis: Calcd. for $C_{19}H_{26}N_2O_3$:

C, 68.86; H, 7.90; N, 8.45.

Found: C, 68.57; H, 7.87; N, 8.14.

EXAMPLE 71-Methyl-5-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]-1,2,3-thiadiazole 2,2-Dioxide

The diamines of Example 2 (3.3 g, 11.37 mmol) are dissolved in 180 ml dry pyridine and are treated with sulfamide (1.36 g, 14.22 mmol). The reaction mixture is warmed to reflux and is refluxed for 15 hours. The reaction is cooled to room temperature and is diluted with 500 ml of ethyl ether and is washed with 5 x 100 ml H_2O , 5 x 100 ml 1 normal HCl and 2 x 200 ml water. The organics are dried over anhydrous sodium sulfate, filtered and concentrated in vacuo. The crude residue is chromatographed on SiO_2 with 50% ethyl acetate/hexanes as the eluent. The appropriate fractions are collected, and concentrated to yield 1.35 g (33.7%) of the cyclic sulfamides as an orange solid.

MP: 55.57°C.

1H NMR (300 MHz, $CDCl_3$): delta 6.85-6.7 (m, 3H), 4.7-4.5 (m, 1.5H), 4.3-4.1 (m, 1.5H), 3.8 (bs, 3H), 3.65 (m, 1H), 3.3 (m, 1H), 2.51 (bs, 3H), 2.6-1.0 (m, 10H).

HRMS 352.1457 (M+) calcd. for $C_{17}H_{24}N_2O_4S$, 352.1466.

-39-

EXAMPLE 81-Methyl-5-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]-2-Imidazolidinethione

The diamines of Example 2 (2.3 g, 7.93 mmol) are dissolved in 65 ml of tetrahydrofuran and treated with N,N-thiocarbonyl imidazole (1.76 g, 9.91 mmol). The reaction is stirred for 41 hours at room temperature. The reaction is worked up by dilution with 250 ml of ethyl ether. The collected organics are washed with 1 x 80 ml water, 2 x 100 ml .5N NaOH solution, 2 x 100 ml .5N HCl solution, 1 x 100 ml water. The organics are dried over anhydrous magnesium sulfate, filtered and concentrated in vacuo. The crude residue is purified by flash silica gel chromatography with 50% ethyl acetate/hexanes as the eluent. The appropriate fractions are collected and concentrated to yield 1.70 g (65.5%) a 7:3 endo:exo mixture of the thioimidazolidinone as a light yellow solid.

MP: 149-150.5°C.

¹H NMR (250 MHz, CDCl₃): delta 6.9-6.7 (m, 3H), 6.15 (bs, 1H), 4.7 (m, 1H), 4.6 (m, .7H), 4.2 (m, .3H), 3.9 (m, 3H), 3.86 (s, 2.1H), 3.83 (s, .9H), 3.42 (m, 1H), 2.93 (bs, 3H), 2.6-1.1 (m, 10H).

HRMS 332.1559 (M+) Calcd. for C₁₈H₂₄N₂SO₂ 332.1583.

25

30

-40-

EXAMPLE 9

alpha-N-Methylamino-3-(Exo-Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxybenzene acetonitrile

3-(exo-bicyclo[2.2.1]hept-2-yloxy)-4-methoxybenz-
aldehyde (4.22 g, 17.0 mmol) is dissolved in 60 ml of
ethanol and is treated with methylamine hydrochloride
(1.40 g, 21.27 mmol) and sodium cyanide (1.04 g, 21.27
mmol). To the reaction is added 15 ml of water to make
the mixture homogeneous. The reaction is stirred at room
temperature for 72 hours. The reaction is worked up by
dilution with 250 ml of ethyl ether and is washed with
3 x 100 ml water and 1 x 100 ml saturated sodium
chloride solution. The organics are dried over
magnesium sulfate, filtered and concentrated in vacuo to
yield 4.45 g (91%) of the amino-nitrile as a clear light
yellow oil.

¹H NMR (300 MHz, CDCl₃): delta 7.1-6.8 (m, 3H),
4.7 (bs, 1H), 4.25 (m, 1H), 3.88 (s, 3H), 2.56 (s, 3H),
2.5-1.0 (m, 10H).

-41-

EXAMPLE 102-Methylamino-2-[3-(Exo-bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]ethylamine

5 The exo-bicyclo[2.2.1]hept-2-yloxy-aminonitrile of
Exmple 9 (4.45 g, 15.56 mmol) is dissolved in 100 ml of
dry toluene and is added to a -78°C. solution of Dibal-H
(70.01 mmol) in 300 ml toluene. The reaction is stirred
at -78°C. for 4 hours. The cooling bath is removed and
the reaction is quenched slowly dropwise with 100 ml of
10 a saturated sodium/potassium tartrate solution. The
reaction is warmed slowly to room temperature and is
diluted with 500 ml of ethyl acetate. The layers are
separated and the aqueous layer is saturated with sodium
chloride and extracted 1 x 100 ml methylene chloride.
15 The collected organics are washed with 1 x 100 ml
saturated sodium/potassium tartrate solution and 2 x 100
ml saturated sodium chloride solution. The organics are
dried over potassium carbonate filtered and concentrated
in vacuo to yield 4.0 g (90%) of the exo-bicyclo[2.2.1]-
20 hept-2-yloxy-diamine as a clear viscous oil.

¹H NMR (300 MHz, CDCl₃): delta 6.83 (m, 3H), 4.22
(bd, 1H), 3.83 (s, 3H), 3.38 (m, 1H), 2.84 (m, 2H), 2.3
(s, 3H), 2.5-1.0 (m, 10H).

25

30

-42-

EXAMPLE 111-Methyl-5-[3-Exo-bicyclo[2.2.1]hept-2-yloxy)-
4-Methoxyphenyl]-2-Imidazolidinone

5 The exo-bicyclo[2.2.1]hept-2-yloxy-diamine of
Example 10 (4.0 g, 13.79 mmol) is dissolved in 170 ml of
tetrahydrofuran and is treated with N,N-carbonyldi-
imidazole (2.8 g, 17.24 mmol). The reaction is stirred
for 40 hours at room temperature. The reaction mixture
is worked up by dilution with 300 ml of ethyl ether and
10 is washed with 1 x 100 ml of water, 1 x 100 ml of .5N
sodium hydroxide solution, 1 x 100 ml .5N HCl solution
and 1 x 100 ml H₂O. The organics are dried over sodium
sulfate, filtered and concentrated in vacuo. The
residue is purified on silica gel flash chromatography
15 with 50% ethyl acetate/hexanes as the eluent. The
appropriate fractions are collected and concentrated in
vacuo to yield 1.60 g (36.7%) of the imidazolidinone as
a white powder. MP 148-151°C.

20 ¹H NMR (300 MHz, CDCl₃): delta 6.8 (m, 3H), 5.3
(3, 1H), 4.4 (m, 1H), 4.2 (bd, 1H), 3.80 (s, 3H), 3.68
(dd, J=11.5Hz, J=8Hz), 3.21 (dd, J=11.5Hz, J=8.1Hz), 2.6
(s, 3H), 2.5-1.0 (m, 10Hz).

25 ¹³C NMR (63 MHz, CDCl₃): delta 163.12, 150.03,
147.63, 131.67, 119.43, 119.35, 112.92, 112.82, 112.02,
81.04, 62.74, 56.0, 47.46, 41.0, 39.79, 35.33, 35.23,
28.63, 28.25, 24.15 (20 lines).

HRMS 316.1816 (M+) Calcd. for C₁₈H₂₄N₂O₃ 316.1787.

30

-43-

EXAMPLE 12

alpha-N-Methylamino-3-(Endo-Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxybenzene Acetonitrile

The aminonitrile is prepared from the aldehyde (Preparation AA) (3.46 g, 14.06 mmol) according to the procedure of Example 1 to produce 3.8 g (95%) of the product.

¹H NMR (300 MHz, CDCl₃): delta 7.0-6.75 (m, 3H), 4.62 (bs, 1H), 4.7-4.5 (m, 1H), 3.82 (s, 3H), 2.65 (m, 1H), 2.6 (bs, 3H), 2.3 (m, 1H), 2.05 (m, 2H), 2.2-1.1 (m, 6H).

EXAMPLE 13

2-Methylamino-2-[3-[Endo-Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]ethylamine

The diamine is prepared according to the procedure of Example 2 from the aminonitrile (3.8 g, 13.28 mmol) of Example 12 to produce 3.9 g (~100%) of the product.

¹H NMR (300 MHz, CDCl₃): delta 6.9-6.7 (m, 3H), 4.7 (m, 1H), 3.91 (s, 3H), 3.96 (m, 1H), 2.96 (m, 2H), 2.6 (m, 1H), 2.3 (bs, 3H), 2.3-1.0 (m, 9H).

-44-

EXAMPLE 141-Methyl-5-[3-(Endo-Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]-2-Imidazolidinone

The imidazolidinone is prepared from the diamine of Example 13 (3.9 g, 13.44 mmol) according to the method of Example 4 to afford 1.6 g (38%) of the product as a white solid. MP: 148-149.5°C.

¹H NMR (300 MHz, CDCl₃): delta 6.85-6.65 (m, 3H), 4.6 (m, 1H), 4.45 (m, 1H), 3.83 (s, 3H), 3.68 (m, 1H), 3.2 (m, 1H), 2.6 (s, 3H), 2.55 (m, 1H), 2.3-1.1 (m, 9H).

¹³C NMR (63 MHz, CDCl₃): delta 163.1, 149.9, 148.8, 131.7, 119.3, 112.4, 112.26, 112.11, 79.01, 62.82, 62.84, 56.21, 47.5, 40.61, 37.26, 37.23, 37.13, 36.77, 29.42, 28.76, 28.73, 27.32, 29.74 (23 lines).

Some doubling due to remote diastereomers.

-45-

EXAMPLE 15

alpha-N-Methylamino-3-(Exo-Bicyclo[2.2.2]oct-2-yloxy)-4-Methoxybenzene Acetonitrile

The bicyclo[2.2.2]octylether of Preparation BB
5 (3.30 g, 12.7 mmol) is dissolved in 50 ml of ethanol and
to this is added methylamine hydrochloride (1.28 g, 19
mmol) and sodium cyanide (930 mg, 19 mmol). To this
suspension is added dropwise water until the reaction
mixture becomes clear. The reaction is stirred for 15
10 hours and is then treated with additional methylamine
hydrochloride (320 mg, 4.75 mmol) and sodium cyanide
(232 mg, 4.75 mmol). The reaction is stirred for 24
hours at room temperature. The reaction is diluted with
250 ml H₂O and 250 ml ethyl ether. The resulting
15 aqueous layer was re-extracted with 100 ml ether and the
combined organics are washed with 1 x 200 ml 1N NaOH
solution, 2 x 200 ml water and 1 x 100 ml brine. The
organics are dried over magnesium sulfate, filtered and
concentrated in vacuo to yield 3.80 gram (✓100%) of the
20 aminonitrile as a viscous oil. It was used in Example
16 without further purification.

25

30

-46-

EXAMPLE 162-Methylamino-2-[3-(Exo-Bicyclo[2.2.1]oct-2-yloxy)-4-Methoxyphenyl]ethylamine

To a solution of Dibal-H (64.5 mmol) in 140 ml of
toluene at -78°C. is added dropwise the cyanoamine (3.80
g, 12.7 mmol) of Example 15 over a 1 hour period as a
solution in 50 ml toluene. After the addition is
complete the reaction is stirred for an additional 1.5
hours at -78°C. The reaction mixture is quenched at
-78°C. with a saturated solution of sodium potassium
tartrate (5 ml) and allowed to warm slowly to room
temperature. The reaction is then treated dropwise with
an additional 100 ml of saturated sodium potassium
tartrate solution and is stirred for 15 hours. The
suspension is diluted with 100 ml ethyl acetate. The
aqueous layer is saturated with sodium chloride and
extracted 2 x 100 ml ethyl acetate. The collected
organics are washed with brine dried over magnesium
sulfate, filtered and concentrated in vacuo to yield
3.14 g (81%) of the aryldiamine as a viscous oil. It
was used as is in the procedure of Example 17.

25

30

-47-

EXAMPLE 171-Methyl-5-[3-Exo-Bicyclo[2.2.1]oct-2-yloxy)-
Methoxyphenyl]-2-Imidazolidinone

5 The aryldiamine (3.14 g, 10.3 mmol) of Example 16 is dissolved in 100 ml of dry tetrahydrofuran is treated with N,N-carbonyldiimidazole (2.09 g, 12.9 mmol) and is stirred at room temperature for 90 hours. The reaction mixture is diluted with 200 ml of ethyl ether and is washed with 1 x 50 ml 2.5% HCl solution, 1 x 50 ml water, 1 x 50 ml 1N NaOH solution, 1 x 50 ml H₂O and 1 x 50 ml brine. The organics are dried over MgSO₄, filtered and concentrated in vacuo. The residue is chromatographed on SiO₂ (32-64) with 20% hexane/80% ethyl acetate as the eluent. The appropriate fractions are collected and concentrated in vacuo to yield 1.10 g (32%) of the imidazolidinone as a white solid. MP: 142-146°C.

20 ¹H NMR (300 MHz, CDCl₃): delta 6.85-6.65 (m, 3H), 5.4 (bs, 1H), 4.45-4.3 (m, 2H), 3.8 (s, 3H), 3.65 (m, 1H), 3.2 (m, 1H), 2.55 (bs, 3H), 2.1-1.2 (m, 12H).

25 ¹³C NMR (75.6 MHz, CDCl₃): delta 163.2, 163.14, 150.46, 150.43, 147.9, 131.7, 119.68, 119.63, 113.54, 113.4, 112.3, 76.24, 76.18, 62.86, 62.80, 62.76, 56.22, 47.61, 34.86, 28.80, 28.41, 25.37, 24.62, 23.42, 19.21 (25 lines). Some doubling due to remote diastereomers.

HRMS 330.1943 (M+) calculated for C₁₉H₂₆N₂O₃
330.1924 (M+) calculated to C₁₉H₂₆N₂O₃ 330.1924.

-48-

EXAMPLE 18alpha-N-Methylamino-3-(Exo-Bicyclo[3.2.1.]oct-2-yloxy)-4-Methoxybenzene Acetonitrile

5 The aryl ether aldehyde of Preparation CC (1.68 g, 6.46 mmol) is dissolved in 50 ml of ethanol and is treated with sodium cyanide (.379 g, 7.75 mmol) and methylamine hydrochloride (.519 g, 7.75 mmol). The reaction is stirred for 24 hours at room temperature. The reaction is made basic with saturated NaHCO_3 solution and is diluted with 50 ml of water. The aqueous layer is extracted 3 x 30 ml ethyl ether. The combined organics are washed with 2 x 30 ml water and 2 x 30 ml brine, dried over Na_2SO_4 , filtered and concentrated in vacuo to yield 1.65 g (89.5%) of the aminonitrile as a clear orange oil.

10

15

^1H NMR (300 MHz, CDCl_3): delta 7.05-6.75 (m, 3H), 4.65 (bs, 1H), 4.45 (m, 1H), 3.83 (s, 3H), 2.55 (bs, 3H), 2.3 (m, 1H), 2.05 (m, 1H), 1.8-1.3 (m, 10H).

20

25

30

-49-

EXAMPLE 192-Methylamino-2-[3-(Exo-Bicyclo[3.2.1]oct-2-yloxy)-4-Methoxyphenyl]ethylamine

5 A solution of diisobutyl aluminum hydride (20.25 ml of a 1.5 molar solution in toluene, .028 mol) in 200 ml of dry toluene is cooled to -78°C. and is treated dropwise with a solution of the aminonitrile of Example 18 (1.65 g, 5.76 mmol) in 25 ml of toluene. The addition is completed in 15 hours and the reaction is stirred at -78°C. for 2 hours. The reaction is warmed to 0°C. and is quenched with 50 ml of a saturated sodium potassium tartrate solution. The organic layer is separated and the aqueous layer is extracted 3 x 30 ml ether. The combined organics are washed with 3 x 30 ml dilute sodium potassium tartrate solution, 3 x 30 ml brine. The organics are dried over NaSO₄, filtered and concentrated in vacuo to yield .95 g (54.5%) of the diamine as a clear yellow oil.

20 ¹H NMR (300 MHz, CDCl₃): delta 6.9-6.75 (m, 3H), 4.45 (m, 1H), 3.8 (s, 3H), 3.4 (m, 1H), 2.85 (m, 2H), 2.32 (s, 3H), 2.2-1.4 (m, 12H).

25

30

-50-

EXAMPLE 201-Methyl-5-[3-(Exo-Bicyclo[2.2.1]oct-2-yloxy)-
4-Methoxyphenyl]-2-Imidazolidinone

5 The aryl diamine of Example 19 (.95 g, 3.14 mmol)
is dissolved in 30 ml of dry tetrahydrofuran and is
treated with N,N-carbonyldiimidazole (.76 g, 4.71 mmol).
The reaction is stirred for 18 hours at room temperature
and is then diluted with 30 ml water and 30 ml ethyl
acetate. The aqueous layer is separated and extracted
10 2 x 20 ml ethyl acetate. The combined organics are
washed with 2 x 20 ml in NaOH solution, 2 x 20 ml 1N
HCl, 2 x 20 ml water and 2 x 20 ml brine. The organic
layer is dried over Na₂SO₄, filtered and concentrated in
vacuo to yield a white slurry. Purification by
15 titration with 3 x 50 ml ether to yield 214 mg (20.6%)
of the imidazolidinone as a white solid. MP:
145-147°C.

¹H NMR (300 MHz, CDCl₃): delta 6.9-6.7 (m, 3H),
4.5-4.3 (m, 2H), 3.83 (s, 3H), 3.7 (m, 1H), 3.24 (m,
20 1H), 2.62 (s, 3H), 2.3 (m, 1H), 2.05 (m, 1H), 1.8-1.3
(m, 10H).

HRMS 330.1962 (M+) Calcd for C₁₉H₂₆N₂O₃ 330.1943.

25

30

-51-

EXAMPLE 21

5-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxy-phenyl]-2,4-Imidazolidinedione

3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxybenz-
5 aldehyde (20.0 g, 81.3 mmol), sodium cyanide (8.0 g,
162.6 mmol) and ammonium carbonate (32.0 g, 333.3 mmol)
are dissolved in 100 ml ethanol and 100 ml water and
refluxed for 4 hours. The reaction is cooled, neutra-
lized with 1N HCl solution and the product extracted 2x
10 ethyl acetate. The collected organics are washed with
water, brine and concentrated in vacuo. The solid
is redissolved in ethyl acetate, dried over Na₂SO₄,
concentrated in vacuo and the resulting crude oil
trituated from ether to produce 18.3 g (71%) of the
15 hydantoin as a crystalline material. This material is a
7:3 endo/exo mixture of bicycloalkyl isomers.

¹H NMR (300 MHz, DMSO): delta 7.4-6.8 (m, 3H), 5.3
(m, 1H), 4.7 (m, .7H), 4.3 (m, .3H), 3.9 (s, 3H),
2.6-1.0 (m, 10H).

20 HRMS 316.1433 Calcd. for C₁₇H₂₀N₂O₄ 316.1450.

25

30

-52-

EXAMPLE 224-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxy-phenyl]-2,4-Imidazolidinedione

5 The 2,4-imidazolidinedione of Example 21 (3.0 g, 9.5 mmol) is dissolved in 40 ml THF and treated with 19 ml of a 1 molar solution of lithium aluminum hydride in THF. The reaction is refluxed for 48 hours, cooled and quenched with 10 ml saturated Na-K tartrate solution. The reaction is extracted 2x ethyl acetate, dried over
10 MgSO₄, filtered, concentrated and flashed on SiO₂ with hexane/ethyl acetate (1:1) → 20% EtOH/hexanes as the eluent. Obtained is 610 mg (21.2%) of the product as a crystalline solid. This material is 75% endo isomer.

MP = 146-148°C.

15 ¹H NMR (300 MHz, CDCl₃): delta 6.8 (m, 3H), 5.2 (m, 2H), 4.8 (m, 1H), 4.6 (m, .7H), 4.2 (m, .3H), 3.9 (bs, 3H), 3.8 (m, 1H), 3.3 (m, 1H), 2.7-1.0 (m, 10H).

HRMS 302.1641 Calcd. for C₁₇H₂₂N₂O₃ 302.1630.

Analysis:

20 Calcd.: C, 67.52; H, 7.34; N, 9.27.

Found: C, 67.37; H, 7.30; N, 9.19.

25

30

-53-

EXAMPLE 23

alpha-N-Methylamine-3-(Endo-Tricyclo[3.3.1.1^{3,7}]-
dec-2-yloxy)-4-Methoxybenzene Acetonitrile

The 2-adamantyl isovanillin ether of Preparation DD
(5.29 g, 18.5 mmol) is dissolved in 150 ml of ethanol
and is treated with sodium cyanide (1.36 g, 27.74 mmol)
and methylamine hydrochloride (1.83 g, 27.74 mmol). To
this is added 20 ml of water to homogenize the reaction
mixture. The reaction is stirred for 48 hours at room
temperature and is worked up by dilution with 300 ml
ethyl ether and washing with 3 x 100 ml water. The
organics are dried over N₂SO₄, filtered and concentrated
in vacuo to yield 5.28 g (88%) of the aminonitrile as a
viscous oil.

¹H NMR (300 MHz, CDCl₃): delta 7.0-6.75 (m, 3H),
4.6 (bs, 1H), 4.35 (bs, 1H), 3.8 (s, 3H), 2.5 (bs, 3H),
2.3-1.4 (m, 14H).

-54-

EXAMPLE 242-Methylamino-2-[3-(Endo-Tricyclo[3.3.1.1^{3,7}]-
dec-2-yloxy)-4-Methoxyphenyl]ethylamine

The aminonitrile of Example 23 (5.95 g, 18.25 mmol) is dissolved in 200 ml of dry toluene and is added to a -78°C. solution of Dibal-H (91.25 mmol) in 270 ml of dry toluene. The reaction is stirred at -78°C. for 4 hours. The cooling bath is removed and the reaction is quenched slowly dropwise with 150 ml of a saturated sodium/potassium tartrate solution. The reaction is warmed slowly to room temperature and is diluted with 500 ml of ethyl acetate. The layers are separated and the aqueous is saturated with sodium chloride and extracted 1 x 100 ml methylene chloride. The collected organics are washed with 1 x 100 ml saturated sodium/potassium tartrate solution and 2 x 100 ml brine. The organics are dried over K₂CO₃ filtered and concentrated in vacuo to yield 6.0 g (~100%) of the adamantyl isovanillin diamine as a clear viscous oil.

¹H NMR (300 MHz, CDCl₃): delta 6.9-6.7 (m, 3H), 4.4 (bs, 1H), 3.85 (s, 3H), 3.35 (m, 1H), 2.8 (m, 2H), 2.3 (s, 3H), 2.2-1.4 (m, 14H).

-55-

EXAMPLE 25

1-Methyl-5-[3-(Endo-Tricyclo[3.3.1.1^{3,7}]dec-2-yloxy)-4-Methoxyphenyl-2-Imidazolidinone

5 The diamine of Example 24 (6.0 g, 18.2 mmol) is dissolved in 180 ml of tetrahydrofuran and is treated with N,N-carbonyldiimidazole and is stirred at room temperature for 24 hours. The reaction mixture is diluted with 250 ml of ethyl ether and 200 ml of water. The organics are washed with 1 x 100 ml .5N NaOH
10 solution, 1 x 100 ml .5N HCl solution, 1 x water, dried over Na₂SO₄, filtered and concentrated in vacuo. The residue is chromatographed on SiO₂ with 50% ethyl acetate/hexanes as the eluent. The appropriate fractions are collected and concentrated in vacuo to
15 yield 1.85 g (28%) of the adamantyl imidazolidinone as a white crystalline solid. MP = 180.5-181°C.

¹H NMR (300 MHz, CDCl₃): delta 6.8 (bs, 2H), 5.4 (bs, 1H), 4.45 (m, 2H), 3.82 (s, 3H), 3.66 (m, 1H), 3.18 (m, 1H), 2.57 (s, 3H), 2.4-1.5 (m, 14H).

20 HRMS 356.2115 (M+) calcd. for C₂₁H₂₈N₂O₃ 356.2100.

25

30

-56-

EXAMPLE 26alpha-N-Ethylamino-3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxybenzene Acetonitrile

5 The 3-(bicyclo[2.2.1]hept-2-yloxy)-4-methoxy-
benzaldehydes of Example 1 (2.3 g, 9.35 mmol) are
dissolved in 60 ml of ethanol and are treated with
sodium cyanide (0.60 g, 12.2 mmol) and ethyl amine
hydrochloride (1.0 g, 12.2 mmol). The solution is made
homogenous with 4 ml of water and is stirred for 48
10 hours at room temperature. The reaction mixture is
diluted with 250 ml of ethyl ether and 20 ml of
saturated NaHCO₃ solution. The organics are washed with
3 x 100 ml water, 3 x 100 ml brine, dried with Na₂SO₄,
filtered and concentrated in vacuo to yield 2.3 g (82%)
15 of the ethyl aminonitrile as a viscous yellow oil. This
material represents a 7:3 mixture of endo/exo isomers at
the bicycloalkyl ether linkage.

¹H NMR (300 MHz, CDCl₃): delta 6.9-6.7 (m, 3H),
4.6 (bs, 1H), 4.55 (m, 7H), 4.1 (m, 3H); 3.7 (s, 3H),
20 2.8-1.0 (m, 15H).

25

30

-57-

EXAMPLE 272-Ethylamino-2-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]ethylamine

The ethylamino nitrile of Example 26 (2.3 g, 7.7 mmol) is dissolved in 70 ml of dry toluene and is added to a -78°C. solution of Dibal-H (38.5 mmol) in 100 ml toluene. The reaction is stirred for 2 hours at 78°C. and is warmed slowly to -40°C. where it is quenched slowly with 60 ml of a saturated sodium potassium tartrate solution. The reaction is then warmed to room temperature and is diluted with 200 ml of ethyl ether. The layers are separated and the aqueous is re-extracted with 100 ml ethyl acetate. The collected organics are washed with 3 x Na K tartrate solution, 2 x 100 ml water, 3 x 100 ml brine, dried over Na₂SO₄, filtered and concentrated in vacuo to yield 1.9 g (81.2%) of the diamine as a brown oil. This material represents a 7:3 endo/exo mixture of isomers at the bicycloalkyl ether linkage.

¹H NMR (300 MHz, CDCl₃): delta 6.85-6.7 (m, 3H), 4.6 (m, .7H), 4.2 (m, .3H), 3.8 (bs, 3H), 3.65 (bs, 1H), 3.5 (m, 1H), 2.8 (m, 2H), 2.7-1.0 (m, 15H).

-58-

EXAMPLE 281-Ethyl-5-[3-(Bicyclo[2.2.1]hept-2-yloxy)-
4-Methoxyphenyl]-2-Imidazolidinone

5 The ethyl-diamine of Example 27 (1.8 g, 5.9 mmol)
is dissolved in 20 ml of dry tetrahydrofuran, is treated
with N,N-carbonyldiimidazole (1.4 g, 8.8 mmol) and is
stirred at room temperature for 24 hours. The reaction
mixture is diluted with 100 ml of water and extracted
2 x 100 ml ethyl acetate. The collected organics are
10 washed 2 x 50 ml .5 normal NaOH solution, 2 x 50 ml .5
normal HCl solution, 1 x 50 ml water and 2 x 50 ml
brine. The organics are dried over Na₂SO₄ filtered and
concentrated in vacuo. The resulting oil is titerated
with ether, washed with ether and dried in vacuo to
15 yield 579 mg (29.7%) of the imidazolidinone as a white
solid. This material represents a 7:3 endo/exo mixture
of isomers at the bicycloalkyl ether linkage. MP =
149-153°C.

¹H NMR (300 MHz, CDCl₃): delta 6.8-6.7 (m, 3H),
20 5.4-5.1 (bs, 1H), 4.55 (m, 1.7H), 4.15 (m, .3H), 3.8 (s,
2.1H), 3.77 (s, .7H), 3.6 (m, 1H), 3.4 (m, 1H), 3.2 (m,
1H), 2.72 (m, 1H), 2.54 (m, .7H), 2.45 (m, .3H), 2.27
(m, .3H), 2.23 (m, .7H), 2.0 (m, 2H), 1.8-1.1 (m, 6H),
.98 (bt, 3H, J=7Hz).

25 HRMS 330.1951(+) calcd. for C₁₉H₂₆N₂O₃ 330.1943.

30

-59-

EXAMPLE 29alpha-N-Allylamino-3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxybenzene Acetonitrile

5 The 3-(bicyclo[2.2.1]hept-2-yloxy)-4-methoxybenz-
aldehydes of Example 1 (2.3 g, 9.35 mmol) are dissolved
in 60 ml of ethanol and treated with sodium cyanide
(0.60 g, 12.2 mmol), allyl amine (.9 ml, 12.2 mmol) and
1.02 ml of concentrated HCl solution. The reaction
mixture is stirred for 48 hours at room temperature.
10 The reaction is diluted with 250 ml ethyl ether and 20
ml of saturated NaHCO₃ solution. The organics are
washed with 3 x 100 ml water, 3 x 100 ml brine, dried
over sodium sulfate, filtered and concentrated in vacuo
to yield 2.7 g (92%) of the allyl cyanoamine as a
15 viscous yellow oil. This material represents a 7:3
endo/exo mixture of isomers at the bicycloalkyl ether
linkage.
- ¹H NMR (300 MHz, CDCl₃): delta 6.9-6.6 (m, 3H),
5.75 (m, 1H), 5.18 (bd, 1H, J=15Hz), 5.07 (bd, 1H,
20 J=9Hz), 4.6 (bs, 1H), 4.5 (m, .7H), 4.07 (m, .3H), 3.7
(s, 3H), 3.4-3.2 (m, 2H), 2.6-1.0 (m, 10H).

25

30

-60-

EXAMPLE 302-Allylamino-2-[3-Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]ethylamine

The allyl aminonitrile mixture of Example 29 (2.7 g, 8.6 mmol) is dissolved in 80 ml of dry toluene and is added dropwise to a -78°C. solution of Dibal-H (43 mmol) in 110 ml of dry toluene. The reaction is stirred to 2 hours at -78°C. and is warmed slowly to -40°C. where it is quenched slowly with 70 ml of saturated sodium potassium tartrate solution. The reaction is then warmed to room temperature and is diluted with 250 ml ethyl ether. The layers are separated and the aqueous is re-extracted with 150 ml ethyl acetate. The collected organics are washed with 3 x 50 ml Na K tartrate solution, 2 x 100 ml water, 3 x 100 ml brine, dried over Na₂SO₄, filtered and concentrated in vacuo to yield 2.4 g (88.3%) of the allyl diamine as a viscous pale brown oil. This material represents a 7:3 endo/exo mixture of isomer at the bicycloalkyl ether linkage.

¹H NMR (300 MHz, CDCl₃): delta 6.85-6.7 (m, 3H), 5.8 (m, 1H), 5.08 (bd, J=15Hz), 5.01 (bd, J=9Hz), 3.73 (s, 3H), 3.5 (m, 1H), 3.1 (m, 2H), 2.75 (m, 2H), 2.6-1.0 (m, 10H).

-61-

EXAMPLE 311-Allyl-5-[3-(Bicyclo[2.2.1]hept-2-yloxy)-
4-Methoxyphenyl]-2-Imidazolidinone

5 The allyldiamine mixture of Example 30 (2.3 g, 7.3 mmol) is dissolved in 30 ml of dry tetrahydrofuran, is treated with N,N-carbonyldiimidazole, and is stirred at room temperature for 24 hours. The reaction mixture is diluted with 100 ml of water and extracted 2 x 100 ml ethyl acetate. The collected organics are washed 2 x 50 ml .5 normal NaOH solution, 2 x 50 ml. 5 normal HCl solution, 1 x 50 ml water, and 2 x 50 ml brine. The organics are dried over Na₂SO₄ filtered and concentrated in vacuo. The resulting oil is titerated with ether, washed with ether and dried in vacuo to yield 739 mg (29.6%) of the allyl imidazolidinone as a white solid. This material represents a 7:3 endo/exo mixture of isomers at the bicycloalkyl ether linkage. MP = 110-113°C.

20 ¹H NMR (300 MHz, CDCl₃): delta 6.8-6.65 (m, 3H), 5.6 (m, 1H), 5.06 (bd, 1H, J=9Hz), 4.98 (bd, 1H, J=15Hz), 4.6 (m, 1.7H), 4.1 (m, 1.3H), 3.8 (s, 3H), 3.7 (m, 1H), 3.28 (m, 1H), 3.1 (dd, 1H, J=12Hz, H=8Hz), 2.57 (m, .7H), 2.48 (m, .3H), 2.3-2.2 (m, 1H), 2.1-1.1 (m, 8H).

25 HRMS 342.1971(+) Calcd. for C₂₀H₂₆N₂O₃ 342.1943.

-62-

EXAMPLE 32alpha-N-Phenylethylamino-3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxybenzene Acetonitrile

5 The norbornyl isovanillin aldehyde (2.3 g, 9.35
mmol) is dissolved in 60 ml of ethanol and is treated
with sodium cyanide (0.60 g, 12.2 mmol), phenethylamine
(1.5 ml, 12.2 mmol), and 1.02 ml of concentrated
hydrochloric acid solution. The reaction is diluted
10 with 250 ml of ethyl ether and 20 ml of saturated NaHCO₃
solution. The organics are washed with 3 x 100 ml
water, 3 x 100 ml brine, dried over sodium sulfate,
filtered and concentrated in vacuo to yield 3.6 g
(~100%) of the phenethylaminonitrile as viscous yellow
15 oil. This material represents a 7:3 mixture of isomers
at the bicyclo[2.2.1]hept-2-yl ether linkage.

¹H NMR (300 MHz, CDCl₃): delta 7.4-7.1 (m, 5H),
6.9-6.7 (m, 3H), 4.65 (bs, 1H), 4.55 (m, .7H), 4.1 (m,
.3H), 3.78 (bs, 3H), 3.1-2.5 (m, 5H), 2.3-1.0 (m, 9H).

20

25

30

-63-

EXAMPLE 332-Phenylethylamino-2-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]ethylamine

5 The phenethylaminonitrile mixture of Example 32 (3.6 g, 9.4 mmol) is dissolved in 90 ml of dry toluene and is added dropwise to a -78°C. solution of diisobutylaluminumhydride (47 mmol) in 120 ml of dry toluene. The reaction is stirred for 2 hours at -78°C. and is warmed slowly to -40°C where it is quenched
10 slowly with 70 ml of saturated sodium potassium tartrate solution. The reaction is then warmed to room temperature and diluted with 250 ml ethyl ether. The layers are separated and the aqueous is re-extracted with 200 ml ethyl acetate. The collected organics are
15 washed with 3 x 50 ml Na K tartrate solution, 2 x 100 ml water, 3 x 100 ml brine, dried over Na₂SO₄, filtered and concentrated in vacuo to yield 3.4 g (95%) of the phenethyldiamine as a pale brown viscous oil. This material represents a 7:3 endo/exo mixture of isomers at
20 the bicycloalkyl ether linkage.

¹H NMR (300 MHz, CDCl₃): delta 7.4-7.1 (m, 5H), 6.9-6.7 (m, 3H), 4.6 (m, .7H), 4.2 (m, .3H), 3.8 (bs, 3H), 3.7 (bs, 1H), 3.55 (m, 1H), 3.1-2.5 (m, 7H), 2.5-1.0 (m, 9H).

25

30

-64-

EXAMPLE 341-Phenylethyl-5-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]-2-Imidazolidinone

5 The phenethyldiamine endo/exo mixture of Example 33 (3.3 g, 8.7 mmol) is dissolved in 40 ml of dry tetrahydrofuran, treated with N,N'-carbonyldiimidazole, and is stirred at room temperature for 24 hours. The reaction mixture is diluted with 100 ml of water and extracted 2 x 100 ethyl acetate. The collected organics are washed 2 x 50 ml .5 normal NaOH solution, 2 x 50 ml .5 normal HCl solution, 1 x 50 ml water and 2 x 5 ml brine. The organics are dried over Na₂SO₄, filtered and concentrated in vacuo. The residue is flash chromatographed (32-60 SiO₂ 50% ethyl acetate/hexane → 100% ethyl acetate). The appropriate fractions are concentrated in vacuo and titerated with ether to yield 337 mg (10.6%) of the phenethylimidazolidinone as a white solid. This material represents a 7:3 endo/exo mixture of isomers at the bicycloalkyl ether linkage. MP = 151-154°C.

¹H NMR (300 MHz, CDCl₃): delta 7.4-7.1 (m, 5H), 6.9-6.7 (m, 3H), 4.9 (bs, 1H), 4.6 (m, .7H), 4.45 (m, 1H), 4.2 (m, .3H), 3.83 (bs, 3H), 3.7 (m, 2H), 3.26 (m, 1H), 3.1-2.5 (m, 4H), 2.4-1.1 (m, 9H).

25 HRMS 406.2294 (M+) Calcd. for C₂₅H₃₀N₂O₃ 406.2257.

EXAMPLE 354-[3-(Endo-Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]imidazole Hydrochloride

The 3-(Endo-bicyclo[2.2.1]hept-2-yloxy)-4-methoxy-phenylglyoxal reactant (.30 g, 1.03 mmol) and aqueous CH_2O (3 ml of a 37% aqueous solution) are dissolved in 3 ml of ethanol and to this is added 3 ml of concentrated NH_4OH . The reaction mixture is stirred for 1.5 hours at room temperature and is quenched with 20 ml water. The reaction mixture is extracted 2 x ethyl acetate. The combined organics are mixed and adjusted to pH 2. The aqueous is separated and is basified to pH 9 with 50% NaOH . The aqueous is extracted 3 x ethyl acetate. The organics are washed with water, dried over MgSO_4 , filtered and concentrated in vacuo to yield .2 g of the imidazole as an oil. The oil is dissolved in 1 ml of acetone and is treated with 2 ml of HCl saturated acetone. The solution is treated with ether and the precipitate is collected to yield .12 g (68.4%) of the imidazole as its HCl salt. MP = 201.202°C. (decomp).

^1H NMR (300 MHz, CDCl_3): delta 6.9-6.7 (m, 4H), 6.45 (m, 1H), 4.2 (m, 1H), 4.5 (s, 3H), 2.2 (m, 1H), 1.87 (m, 1H), 1.8-.7 (m, 9H).

HRMS 284.1515(M+) Calcd. for $\text{C}_{17}\text{H}_{20}\text{N}_2\text{O}_3$ 284.1525.

EXAMPLE 36

3-(Indan-2-yloxy)-4-methoxybenzaldehyde (2.25 g, 839 mmol) is converted to the aminonitrile (2.36 g, 91.3%) according to the procedure of Example 2.

^1H NMR (300 MHz, CDCl_3): delta 7.3-7.05 (m, 6H), 6.85 (m, 1H), 5.2 (m, 1H), 4.7 (bs, 1H), 3.8 (s, 3H), 3.45 (dd, 2H, J=13Hz, J=7Hz), 3.25 (dd, 2H, J=13Hz, J=3Hz), 2.6 (s, 3H).

-66-

EXAMPLE 372-Methylamino-2-[3-(Indan-2-yloxy)-4-Methoxyphenyl]ethylamine

5 The aminonitrile of Example 36 (2.25 g, 7.3 mmol) is reduced according to the procedure of Example 3 to produce the diamine (1.5 g, 65.5%) as a clear viscous oil.

¹H NMR (300 MHz, CDCl₃): delta 7.3-7.1 (m, 4H), 6.9 (m, 3H), 5.25 (m, 1H), 3.8 (s, 3H), 3.5 (m, 1H), 3.45 (dd, 2H, J=13Hz, J=7Hz), 3.25 (dd, 2H, J=13Hz, J=3Hz), 2.87 (m, 2H), 2.4 (bs, 3H).

EXAMPLE 381-Methyl-5-[3-(Indan-2-yloxy)-4-Methoxyphenyl]-2-Imidazolidinone

15 The diamine of Example 37 (1.50 g, 4.8 mmol) is converted to the imidazolidinone (126 mg, 37 mmol, 7.75%) according to the procedure of Example 4.

¹H NMR (300 MHz, CDCl₃): delta 7.3-7.2 (m, 4H), 6.9 (m, 3H), 5.3-5.2 (m, 1H), 5.0 (bs, 1H), 4.54 (m, 1H), 3.9 (s, 3H), 3.78 (m, 1H), 3.45 (dd, 2H, J=13Hz, J=7Hz), 3.3 (m, 1H), 3.25 (dd, 2H, J=13Hz, J=3Hz), 2.75 (s, 3H).

25

30

-67-

EXAMPLE 39

alpha-N-Methylamino-3-(Exo-Tricyclo[3.2.1.0^{2,6}]-
dec-8-yloxy)-4-Methoxybenzene Acetonitrile

5 The 3-exo-tricyclo[3.2.1.0^{2,6}]dec-8-yloxy)-4-methoxybenzaldehyde reactant (1.28 g, 4.47 mmol) is dissolved in 60 ml of ethanol and is treated with sodium cyanide (274 mg, 5.6 mmol) and methylamine hydrochloride (370 mg, 5.6 mmol). To this is added 20 ml of water to make the reaction homogeneous. The reaction is stirred
10 for 20 hours at room temperature and is worked up by dilution with 250 ml of ether and is washed with 2 x 100 ml water, 1 x 100 ml pH 7 phosphate buffer, 1 x 100 ml brine, dried over Na₂SO₄, filtered and concentrated in vacuo to yield 1.13 g (78%) of the methylamino nitrile
15 viscous oil.

¹H NMR (300 MHz, CDCl₃): delta 7.0-6.7 (m, 3H), 4.2 (m, 2H), 3.8 (s, 3H), 2.55 (s, 3H), 2.3-.8 (m, 14H).

The following compound is prepared in like manner from the appropriate reactant:

20 alpha-N-methylamino-3-(endo-tricyclo[3.2.1.0^{2,6}]-dec-8-yloxy)-4-methoxybenzene acetonitrile in 89.4% yield:

¹H NMR (300 MHz, CDCl₃): delta 7.1-6.81 (m, 3H), 4.7 (bs, 1H), 4.6 (m, 1H), 3.9 (s, 3H), 2.6 (s, 3H),
25 2.8-1.0 (m, 14H).

-68-

EXAMPLE 40

2-Methylamino-2-[3-(Exo-Tricyclo[3.2.1.0^{2,6}]dec-8-yloxy)-4-Methoxyphenyl]ethylamine

5 The title methylamino nitrile of Example 39 (1.94 g, 5.95 mmol) is dissolved in 60 ml of dry toluene and is added to a -78°C. solution of diisobutylaluminum-hydride (29.75 mmol) as a solution in 150 ml of dry toluene. After the addition is complete the reaction is stirred for 4 hours at -78°C. and is then quenched with 10 65 ml of a saturated solution of sodium potassium tartrate. The reaction is warmed slowly room temperature and diluted with 500 ml of ethyl ether. The organic layer is separated and the remaining aqueous layer is saturated with NaCl and extracted with 2 x 100 15 ml ethylacetate. The collected organics are washed with 1 x 100 ml saturated Na K tartrate, 1 x 100 ml brine, dried over K₂CO₃, filtered and concentrated in vacuo to yield 1.8 g (92%) of the exo isomer of the diamine as a yellow viscous oil.

20 ¹H NMR (300 MHz, CDCl₃): delta 6.85-6.65 (m, 3H), 4.15 (m, 1H), 3.75 (s, 3H), 3.38 (m, 1H), 2.8 (m, 2H), 2.3 (s, 3H), 2.4-.8 (m, 14H).

The endo-isomer is similarly prepared from the endo-methylamino nitrile isomer of Example 39:

25 ¹H NMR (300 MHz, CDCl₃): delta 7.0-6.8 (m, 3H), 5.7 (m, 1H), 3.9 (s, 3H), 3.5 (m, 1H), 3.0-1.0 (m, 19H).

30

-69-

EXAMPLE 411-Methyl-5-[3-(Exo-Tricyclo[3.2.1.0^{2,6}]dec-8-yloxy)-4-Methoxyphenyl]-2-Imidazolidinone

5 The title diamine of Example 40 (1.8 g, 5.45 mmol) is dissolved in 125 ml of tetrahydrofuran and is treated with N,N'-carbonyldimidazole and is stirred at room temperature for 40 hours. The reaction is diluted with 300 ml of ether and washed with 2 x 100 ml water, 1 x .05 normal NaOH, 1 x 0.5 normal HCl, 1 x 100 ml water and 1 x 100 ml brine. The organics are dried over NaSO₄ and concentrated in vacuo. The residue is flashed on SiO₂ (32-64) with 50% ethyl acetate/hexanes as the eluent. The appropriate fractions are concentrated in vacuo to yield 625 mg (32%) of the imidazolidinone (endo isomer) as a white solid. MP = 168-170°C.

Analysis Calcd. for C₂₁H₂₈N₂O₃:

C, 70.76; H, 7.92; N, 7.86.

Found: C, 70.54; H, 7.92; N, 7.97.

¹H NMR (300 MHz, CDCl₃): delta 6.85-6.7 (m, 3H), 5.15 (bs, 1H), 4.4 (m, 1H), 4.12 (m, 1H), 3.81 (s, 3H), 3.68 (m, 1H), 3.22 (m, 1H), 2.6 (s, 3H), 2.3-.8 (m, 14H).

HRMS 356.2155(M⁺) Calcd. for C₂₁H₂₈N₂O₃ 356.2100.

25 The isomeric tricycloalkyl ether is prepared in like manner from the isomeric diamine of Example 40 in 20.7% yield:

MP = 149-152°C.

-70-

^1H NMR (300 MHz, CDCl_3): δ 6.9-6.8 (m, 3H), 5.15 (bs, 1H), 4.65 (m, 1H), 4.5 (m, 1H), 3.9 (s, 3H), 3.75 (m, 1H), 3.25 (m, 1H), 2.7 (s, 3H), 2.8-1.0 (m, 14H).

5 HRMS 356.2120 Calcd. for $\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_3$ 356.2099.

Analysis Calcd. for $\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_3$:

C, 70.76; H, 7.92; N, 7.86.

Found: C, 70.72; H, 7.86; N, 7.79.

EXAMPLE 42

10 α -N-Methylamino-3-(Endo-Tricyclo[3.2.1.0^{2,6}]-dec-4-yloxy)-4-Methoxybenzene Acetonitrile

The aminonitrile is prepared according to the procedure of Example 39 from the corresponding tricyclic isovanillin ether in 95.4% yield.

15 ^1H NMR (300 MHz, CDCl_3): δ 7.0-6.7 (m, 3H), 4.7 (m, 1H), 4.65 (bs, 1H), 3.87, 3.85 (s, 3H), [2-methoxyls], 2.5 (bs, 3H), 2.2-8.5 (m, 14H).

EXAMPLE 43

20 2-Methylamino-2-[3-(Endo-Tricyclo[3.2.1.0^{2,6}]-dec-4-yloxy)-4-Methoxyphenyl]ethylamine

Following the procedure of Example 40, the diamine is prepared from the aminonitrile of Example 42.

25 ^1H NMR (300 MHz, CDCl_3): δ 6.8-6.7 (m, 3H), 4.75 (m, 1H), 3.78 (s, 1H), 3.38 (m, 1H), 2.8 (m, 2H), 2.3 (s, 3H), 2.4-.9 (m, 14H).

-71-

EXAMPLE 44

1-Methyl-5-[3-(Endo-Tricyclo[3.2.1.0^{2,6}]dec-4-yloxy)-4-methoxyphenyl]-2-Imidazolidinone

The imidazolidinone is prepared according to the procedure of Example 41 from the diamine of Example 43 in 7.2% yield.

¹H NMR (300 MHz, CDCl₃): delta 6.8-6.7 (m, 3H), 5.75 (bs, 1H), 4.75 (m, 1H), 4.42 (m, 1H), 3.83 (s, 3H), 3.72 (m, 1H), 3.28 (m, 1H), 2.65 (s, 3H), 2.3-.9 (m, 14H).

¹³C NMR (75.4 MHz, CDCl₃): delta 163.2, 150.8, 147.8, 131.8, 120.0, 114.5, 112.4, 82.5, 62.8, 56.2, 47.6, 46.2, 40.4, 37.7, 31.8, 28.8, 28.5 (17 lines).

Elemental Analysis:

Found: C, 69.74, H, 7.93, N, 7.48

Calcd: C, 70.75, H, 7.91, N, 7.86

-72-

EXAMPLE 45alpha-Formyl-[3-(Exo-Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]Acetonitrile

5 The nitrile of Preparation R (1.2 g, 4.6 mmol) and ethylformate (2 ml) is dissolved in 10 ml benzene and to it is added portionwise sodium hydride [50% in oil] (.39 g, 8.2 mmol). The reaction is warmed to 40°C. for 1.5 hours. The reaction mixture is cooled and diluted with 3 ml ethanol and 20 ml hexanes. The precipitate is
10 collected, suspended in water and acidified with 1N HCl. The product is extracted 3 x 20 ml ethyl acetate and the ethyl acetate extracts are washed with brine, filtered and concentrated in vacuo to yield .62 g (47%) of the formyl compound as a thick yellow oil.

15 ¹H NMR (300 MHz, CDCl₃): delta 7.4-6.7 (m, 4H), 4.25 (m, 1H), 4.25 (m, 1H), 3.8 (s, 3H), 2.6-1.0 (m, 10H).

In like manner, the following alpha-formyl cyanides are prepared from appropriate precursor benzyl cyanide
20 derivatives: alpha-formyl-[3-(bicyclo[2.2.1]hept-2-yloxy)-4-methoxyphenyl]acetonitrile in 68.2% yield as a 7:3 mixture of endo:exo isomers.

¹H NMR (300 MHz, CDCl₃ + CD₃OH): delta 7.2-7.0 (m, 2H), 6.65 (m, 2H), 4.5 (m, .7H), 4.1 (m, .3H), 3.7 (bs, 3H), 2.6-1.0 (m, 10H);
25

alpha-formyl-[3-(indan-2-yloxy)-4-methoxyphenyl]-acetonitrile in 60% yield:

¹H NMR (300 MHz, CDCl₃): delta 7.8 (m, 1H), 7.3 (m, 1H), 7.2-7.1 (m, 4H), 6.9 (m, 3H), 5.15 (m, 1H),
30 3.8 + 3.77 (s, 3H), 3.4-3.1 (m, 4H);

alpha-formyl-[3-(endo-bicyclo[2.2.1]hept-2-yloxy)-4-methoxyphenyl]acetonitrile in 71.8% yield:

¹H NMR (300 MHz, CDCl₃): delta 7.3-7.1 (m, 2H), 6.8 (m, 3H), 4.6 (m, 1H), 3.9 (s, 3H), 2.6-1.0 (m, 10H).

-73-

EXAMPLE 46alpha-Aminomethylene-[3-(Exo-Bicyclo[2.2.1]-hept-2-yloxy)-4-Methoxyphenyl]Acetaldehyde

Water washed Raney-Nickel excess is added to a solution of the title formyl compound of Example 45 (.53 g, 1.81 mol) in 20 ml of ethanol. The reaction mixture is hydrogenated with 40 psi for 7 hours and is filtered through Celite which is subsequently washed with ethanol. The filtrate is concentrated in vacuo to yield .47 g (89.9%) of the imine aldehyde as an oil. This material exists as a mixture of iminealdehyde and enaminoaldehyde.

¹H NMR (300 MHz, CDCl₃): delta 9.6 (d, J=4Hz) + 9.1 (bs) (1H), 7.2-6.6 (m, 4H), 5.1 (bs, 1H), 4.2 (m, 1H), 3.85 (s, 3H), 2.6-1.0 (m, 10H).

The following compounds are similarly prepared from appropriate reactants as tautomeric mixtures:

an approximately 7:3 mixture of endo/exo isomers of alpha-aminomethylene-[3-(endo-bicyclo[2.2.1]hept-2-yloxy)-4-methoxyphenyl] acetaldehyde in 83.3% yield;

alpha-aminomethylene-[3-(endo-bicyclo[2.2.1]hept-2-yloxy)-4-methoxyphenyl]acetaldehyde in 84% yield:

¹H NMR (300 MHz, CDCl₃): delta 9.5 + 9.0 (bs, 1H), 6.9 (m, 3H), 5.4 (bs, 1H), 4.6 (m, 1H), 3.8 (s, 3H), 2.6-1.0 (m, 10H);

alpha-aminomethylene-[3-(indan-2-yloxy)-4-methoxyphenyl]acetaldehyde in 84.6% yield:

¹H NMR (300 MHz, CDCl₃): delta 9.5 + 9.0 (bs, 1H), 7.2-7.1 (m, 4H), 6.9 (m, 3H), 5.3 (bs, 1H), 5.1 (bs, 1H), 3.8 (s, 3H), 3.1 (m, 4H).

-74-

EXAMPLE 475-[3-(Exo-Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]-2-Pyrimidinone

The iminealdehyde of Example 46 (.47 g, 1.63 mmol) is dissolved in 5 ml ethanol and is treated with 1 ml concentrated HCl and urea (.12 g, 1.95 mmol). The reaction is warmed to reflux for 1.5 hours. The reaction mixture is cooled to room temperature, neutralized with aqueous NH_4OH and extracted 6 x 10 ml ethyl acetate. The combined organics are dried over MgSO_4 , filtered and concentrated to dryness to afford .2 g (39.3%) of the pyrimidinone as a crystalline product. MP = 195-196°C.

^1H NMR (300 MHz, CDCl_3): delta 8.4 (m, 2H), 6.9-6.7 (m, 4H), 4.2 (m, 1H), 3.82 (s, 3H), 2.5-.9 (m, 10H).

HRMS 312.1472(+) Calcd. for $\text{C}_{18}\text{H}_{20}\text{N}_2\text{O}_3$ 312.1474.

Similarly the following compounds are prepared from the appropriate precursor enamino aldehydes:

5-[3-(bicyclo[2.2.1]hept-2-yloxy)-4-methoxyphenyl]-2-pyrimidinone in 60.3% yield as a 7:3 endo/exo isomeric mixture.

^1H HMR (300 MHz, CDCl_3): delta 8.4 (m, 2H), 6.9-6.7 (m, 4H), 4.65 (m, .7H), 4.25 (m, .3H), 3.85 (bs, 3H), 2.7-1.1 (m, 10H);

5-[3-(indan-2-yloxy)-4-methoxyphenyl]-1,2-dihydropyrimidinone in 13.9% yield:

^1H NMR (300 MHz, CDCl_3): delta 8.6 (bs, 2H), 7.3-7.0 (m, 7H), 5.35 (m, 1H), 3.75 (s, 3H), 3.4-3.0 (m, 4H);

5-[3-(endo-bicyclo[2.2.1]hept-2-yloxy)-4-methoxyphenyl]-1,2-dihydropyrimidinone in 47% yield:

MP 220°C.

-75-

^1H NMR (300 MHz, CDCl_3): δ 8.5 (bs, 2H), 7.21 (s, 1H), 6.9-6.8 (m, 3H), 4.6 (m, 1H), 3.85 (s, 3H), 2.6-1.0 (m, 10H).

EXAMPLE 48

5 alpha-N-Methylamino-4-Methoxy-3-Nitrobenzene
 Acetonitrile

The aminonitrile is prepared according to the procedure of Example 39 from 4-methoxy-2-nitrobenzaldehyde (25.80 g, 142.4 mmol). The product is isolated by filtration to yield 31.20 g (99%) of product.

10 ^1H NMR (90 MHz, CDCl_3): δ 7.8-7.2 (m, 3H), 5.0 (bs, 1H).

EXAMPLE 49

15 2-Methylamino-2-(4-Methoxy-3-Nitrophenyl)ethylamine

The aminonitrile of Example 48 (15 g, 67.81 mmol) is dissolved in 150 ml of toluene and is cooled to -78°C . To it is added Dibal-H (181 ml, 27.1.2 mmol) as a 1.5 molar solution in toluene. The reaction is stirred at -78°C . for 3 hours, warmed to 0°C . and quenched with 150 ml water. The pH is adjusted to 2 with 6 normal HCl and is washed with 2 x 50 ml ether.

20 The aqueous is pH adjusted to 12 with 25% NaOH solution and is extracted 3 x 100 ml methylene chloride, dried over MgSO_4 , filtered and concentrated in vacuo to yield 11.0 g (72%) of the nitrodiamine as a dark brown oil. This material is used with no further purification.

30

-76-

EXAMPLE 501-Methyl-5-(4-Methoxy-3-Nitrophenyl)-2-Imidazolidinone

The nitroimidazolidinone is prepared by the procedure of Example 40 from the nitrodiamine of Example 49 (11 g, 48.88 mmol) to yield 4.51 g (37%) of the product as a light orange solid.

¹H NMR (90 MHz, CDCl₃): delta 7.7-7.0 (m, 3H), 4.5 (m, 1H), 3.9 (s, 3H), 3.7 (m, 1H), 3.2 (m, 1H), 2.6 (s, 3H).

EXAMPLE 511-Methyl-5-(3-Amino-4-Methoxyphenyl)-2-Imidazolidinone

Platinum oxide (.135 g) is suspended in 1 ml concentrated hydrochloric acid. To this is added to nitroimidazolidinone of Example 50 (4.51 g, 17.95 mmol) as a solution in 30 ml methanol. The volume of the reaction is brought to 200 ml with methanol and is placed on a Parr shaker with 50 psi H₂ pressure for 45 minutes. The catalyst is filtered off and the reaction is concentrated in vacuo to yield an oil which is redissolved in ethyl acetate, washed with 1 normal NaOH solution, dried over MgSO₄, filtered and concentrated in vacuo to yield the crude aniline as an oil which is carried on with no further purification.

¹H NMR (300 MHz, CDCl₃): delta 6.7 (m, 3H), 5.6 (bs, 1H), 4.4 (m, 1H), 3.8 (s, 3H), 3.7 (m, 1H), 3.2 (m, 1H), 2.6 (s, 3H).

-77-

EXAMPLE 521-Methyl-5-[3-(Bicyclo[2.2.1]hept-2-ylamino)-
4-Methoxyphenyl]-2-Imidazolidinone

5 The aniline of Example 51 (3.07 g, 17.95 mmol) is dissolved in 60 ml of glacial acetic acid and to this is added norcamphor (2.37 g, 21.54 mmol). The reaction is cooled to 5°C. and to it is added sodium cyanoboro-
10 hydride (1.36 g, 21.54 mmol). The reaction is poured onto ice and is pH adjusted to 7 with 1 normal NaOH solution. The aqueous is extracted with methylene chloride dried over MgSO_4 , filtered and concentrated in vacuo to afford a brown oil which is flashed on SiO_2 with ethyl acetate as the eluent. The appropriate
15 fractions are combined and concentrated in vacuo to yield a white paste which is tritarted with ether to yield .57 g (10%) of the product as a white crystalline solid. MP = 171-174°C.

^1H NMR (250 MHz, CDCl_3): delta 6.75-6.5 (m, 3H), 4.6 (m, 1H), 4.4 (m, 1H), 3.86 (s, 3H), 3.7 (m, 1H),
20 3.25 (m, 1H), 2.65 (s, 3H), 2.6-2.0 (m, 3H), 1.8-1.2 (m, 6H), .85 (m, 1H).

Elemental Analysis:

Found: C, 68.98, H, 7.40, N, 13.41

Calcd: C, 68.03, H, 7.99, N, 13.37

25 HRMS 315.1948 (M+) Calcd for $\text{C}_{18}\text{H}_{23}\text{N}_3\text{O}_2$ 315.1946.

-78-

EXAMPLE 534-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]pyrazole

The appropriate enamino-aldehyde of Example 46 (0.20 g, 0.697 mM) is dissolved in 5 ml ethanol and treated with .2 ml hydrazine. The reaction mixture is refluxed for 30 minutes, cooled and quenched with water. The product is extracted 2x ethyl acetate and the combined organics are washed, dried over MgSO_4 , filtered and concentrated in vacuo to yield a yellow oil. The crude material is crystallized from ethyl acetate to yield 0.11 g (55.3%) of the pyrazole as a white crystalline solid. This material represents an 7.3 endo to exo mixture at the norbornyl ether linkage.

MP = 180-181°C.

^1H NMR (30 MHz, CDCl_3): delta 7.8 (bs, 2H), 7.1-6.84 (m, 3H), 4.7 (m, .7H), 4.3 (m, .3H), 3.86 (bs, 3H), 2.8-1.0 (m, 10H).

HRMS 284.1531 (M+) Calcd. for $\text{C}_{17}\text{H}_{22}\text{N}_2\text{O}_2$ 284.1525.

-79-

EXAMPLE 546-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]-
imidazo[1,2-a]Pyrimidine

The title enamino-aldehyde of Example 46 (0.20 g, .7 mmol), 2-aminoimidazole sulfate (.27 g, 1 mmol) are dissolved in ethanol, treated with .5 ml, concentrated HCl and refluxed for 1 hour. The reaction is cooled, quenched with water and extracted 3x ethyl acetate. The combined organics are washed with brine, dried over MgSO₄, filtered and concentrated in vacuo. The crude material is crystallized from ethyl acetate to yield 0.20 g (85.3%) of the compound as a beige crystalline solid. This material is 7.3 endo/exo mixture of bicycloalkyl isomers.

MP = 130°C. (Dec.)

¹H NMR (300 MHz, CDCl₃): delta 8.8 (m, 1H), 8.5 (m, 1H), 7.85 (bs, 1H), 7.6 (bs, 1H), 7.2-6.9 (m, 3H), 4.75 (m, 7H), 4.3 (m, 13H), 3.9 (bs, 3H), 2.8-1.1 (m, 10H).

HRMS 335.1610 Calcd. for C₂₀H₂₁O₂N₃ 335.1633.

-80-

EXAMPLE 556-(3-Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]-
pyrazolo[2,3-a]Pyrimidine

5 The title enamino aldehyde of Example 46 (0.20 g,
0.7 mmol) and 3-aminopyrazole (83 mg, 1.0 mmol) are
dissolved in 5 ml of ethanol and treated with .5 ml
concentrated HCl. The reaction mixture is refluxed for
30 minutes, cooled to room temperature, quenched with
water and the product is extracted 2x ethyl acetate.
10 The combined organics are washed with brine, dried over
MgSO₄, filtered and concentrated in vacuo. The crude
material is crystallized from ethyl acetate/hexanes to
produce .180 g (76.7%) of the product as a crystalline
solid comprising endo/exo mixture at the bicycloalkyl
15 residue.

MP = 136-137°C.

¹H NMR (300 MHz, CDCl₃): delta 8.9 (bs, 1H), 8.75
(m, 1H), 8.15 (m, 1H), 7.2-7.0 (m, 3H), 6.8 (m, 1H),
4.75 (m, 7H), 4.3 (m, .3H), 3.9 (bs, 3H), 2.8-1.1 (m,
20 10H).

25

30

-81-

EXAMPLE 56Ethyl 3-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]-2-Carbethoxypropenoate

The aldehyde, 3-(bicyclo[2.2.1]hept-2-yloxy)-4-methoxybenzaldehyde (5.0 g, 0.02 mmole), diethyl malonate (3.21 g, 0.02 mol) and piperidine are dissolved in toluene and refluxed for 15 hours. The reaction is cooled, concentrated in vacuo and the resulting oil taken up in ethyl acetate. The organics are washed with saturated NH_4Cl solution, water, brine, dried, filtered and concentrated in vacuo to afford 6.56 g (84.5%) of diester product as a 7:3 endo/exo isomer mixture.

^1H NMR (300 MHz, CDCl_3): δ 7.6 (m, 1H), 7.2-6.7 (m, 3H), 4.5 (m, .7H), 4.4-4.1 (m, 5H), 3.85 (bs, 3H), 2.6-1.0 (m, 6H).

EXAMPLE 57Ethyl 3-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]-3-Cyanopropanoate

The diester of Example 56 (6.5 g, .0167 mole) and sodium cyanide (.833 g, 0.017 mole) are dissolved in 75 ml of ethanol and stirred at room temperature for 24 hours. The ethanol is removed in vacuo, the resulting solid is partitioned between ethyl acetate and water and the aqueous layer is reextracted with ethyl acetate. The combined organics are washed with water and brine, dried over MgSO_4 , filtered and concentrated in vacuo to afford 4.45 g (77.8%) of the cyano ester as a yellow oil comprising a 7:3 mixture of endo/exo bicycloalkyl isomers.

^1H NMR (300 MHz, CDCl_3): δ 6.9-6.75 (m, 3H), 4.6 (m, 7H), 4.45 (m, 1H), 4.3 (q, 24, $J=5\text{Hz}$), 4.2 (m, .3H), 4.1 (m, 2H), 3.85 (s, 3H), 2.8-1.1 (m, 3H).

-82-

EXAMPLE 584-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxy-phenyl]-2-Pyrrolidone

Platinum oxide (400 mg) is suspended in 50 ml of acetic acid and is activated by hydrogenation at 50 psi for 1 hour. The cyanoester of Example 57 (2.0 g, 5.84 mmol) is added to the PtO₂ suspension as a solution in 50 ml of acetic acid. The reaction mixture is shaken under 50 psi H₂ for 18 hours. The reaction mixture is purged with N₂, concentrated in vacuo and the last traces of acetic acid are azeotroped with toluene in vacuo. The resulting oil is dissolved in 50 ml toluene, treated with 10 ml triethylamine and refluxed for 24 hours. The reaction mixture is then cooled, concentrated in vacuo and the residue dissolved in ethyl acetate. The organics are washed with 1N HCl, water, brine, dried over MgSO₄, filtered and concentrated in vacuo to afford as the alpha-carboethoxy lactam. This material is dissolved in ethanolic NaOH and refluxed until no ester remains by TLC. The reaction mixture is cooled, neutralized with 1N HCl and extracted 3x ethyl acetate. The organics are washed, dried, filtered and concentrated to afford the alpha-carboxylic acid which is thermally decarboxylated at 180°C. to afford 611 mg (34.7%) of the pyrrolidone as a white solid. This material is an 7:3 mixture of endo/exo bicycloalkyl isomers.

MP = 153-156°C.

¹H NMR (300 MHz, CDCl₃): delta 6.9-6.6 (m, 3H), 4.6 (m, .7H), 4.2 (m, .3H), 3.85 (bs, 3H), 3.8 (m, 1H), 3.62 (m, 1H), 3.4 (m, 1H), 2.7 (m, 1H), 2.6 (m, 1H), 2.5 (m, 1H), 2.3 (m, 1H), 2.1-1.0 (m, 8H).

-83-

EXAMPLE 593-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxy-phenyl]-1-Dimethylamino-1-Propen-3-one

1-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-methoxyphenyl]-
5 ethanone (1.5 g, 5.84 mmol) is dissolved in 5 ml
tris-dimethyl aminomethane and refluxed for 30 minutes.
The reaction is cooled, quenched with water and the
product extracted 3x ethyl acetate. The combined
organics are washed with H₂O, brine, dried, filtered and
10 concentrated in vacuo yield 1.4 g (77.3%) of the
enamino-ketone product.

¹H NMR (60 MHz, CDCl₃): delta 7.7 (d, 1H, 12Hz),
7.5-7.2 (m, 2H), 7.0-6.7 (m, 1H), 5.65 (d, 1H, 12Hz),
4.8-4.1 (m, 1H), 3.95 (bs, 3H), 3.0 (s, 6H), 2.9-1.0 (m,
15 10H).

In like manner there is prepared from appropriate
reactants: 3-[(3-indane-2-yloxy)-4-methoxyphenyl]-1-
dimethylamino-1-propen-3-one in 71.8% yield.

¹H NMR (60 MHz, CDCl₃): delta 7.8-6.6 (m, 8H), 6.7
20 (d, 1H, J=12Hz), 3.8 (s, 3H), 3.3 (m, 4H), 2.8 (bs, 6H).

25

30

-84-

EXAMPLE 604-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxy-phenyl]-4-Hydroxy-1,2,3,4-Tetrahydro-2-Pyrimidinone

5 The enamino-ketone of Example 59 (0.9 g, 2.9 mmol) and urea (0.21 g, 3.5 mmol) are dissolved in 10 ml ethanol, treated with 4 ml 1N HCl and refluxed for 1 hour. The reaction is cooled, quenched with water and the product extracted 2x ethyl acetate. The combined
10 organics are washed with brine, dried, filtered and concentrated in vacuo. The crude material is crystallized from CH₂Cl₂/hexanes to yield 0.55 g (57.5%) of the product as beige crystals. This material is 7:3 endo/exo mixture of isomers.

15 MP = 167-169°C. (Dec.)
¹H NMR (300 MHz, CDCl₃): delta 7.7-7.4 (m, 3H), 6.8 (m, 1H), 6.18 (m, 1H), 5.5 (b, 2H), 4.7 (m, .7H), 4.3 (m, .3H), 3.95 (s, 3H), 2.8-1.2 (m, 10H).

HRMS 330.1614 Calcd. for C₁₈H₂₂N₂O₄ 330.1580.

20

25

30

-85-

EXAMPLE 614-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]-
1,2-Dihydro-2-Pyrimidinone

5 The hydroxypyrimidinone of Example 60 (.2 g, .61 mmol) is dissolved in 5 ml ethanol, treated with 2.5 ml 1N HCl and refluxed for 6 hours. The reaction is cooled to room temperature, quenched with water and the product extracted 3x ethyl acetate. The organics are washed, dried, filtered and concentrated in vacuo to yield a
10 crude substance which is crystallized from ethyl acetate to give 60 mg (31.5%) of the pyrimidinone as a crystalline solid. This material is a 7:3 mixture of endo/exo bicycloalkyl isomers.

MP = 220°C.

15 ¹H NMR (300 MHz, CDCl₃): 7.8-7.45 (m, 3H), 6.9 (m, 1H), 6.8 (m, 1H), 5.15 (m, 1H), 4.75 (m, .7H), 4.35 (m, .3H), 3.95 (bs, 3H), 2.8-1.2 (m, 10H).

HRMS 312.1520 Calcd. for C₁₈H₂₀N₂O₃ 312.1474.

20

25

30

-86-

EXAMPLE 625-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxy-phenyl]imidazo[1,2-a]pyrimidine

The enamino-ketone of Example 59 (.4 g, 1.27 mmol) and 2-aminoimidazole sulfate are dissolved in 5 ml ethanol, treated with .5 ml concentrated HCl and refluxed for 1.5 hours. The reaction is cooled, quenched with water, pH adjusted to 9 and the product extracted 2x ethyl acetate. The combined organics are washed, dried, concentrated in vacuo and the crude material crystallized from ether to provide 0.11 g (28.9%) of the product as a beige crystalline solid comprising 7.3: endo/exo mixture of isomers.

MP = 140-141°C.

¹H NMR (300 MHz, CDCl₃): delta 8.6 (d, 1H), 7.81 (bs, 1H), 7.74 (bs, 1H), 7.3-7.0 (m, 3H), 6.8 (m, 1H), 4.63 (m, .7H), 4.21 (m, .3H), 3.95 (bs, 3H), 2.7-1.1 (m, 10H).

HRMS 335.1631 Calcd. for C₂₀H₂₁N₂O₃ 335.1634.

-87-

EXAMPLE 637-[3-(Bicyclo[2.2.1]hept-2-yl)-4-Methoxy-phenyl]pyrazolo[2,3-a]pyrimidine

5 The enamino ketone of Example 59 (0.40 g, 1.27 mmol) and 3-aminopyrazole (0.16 g, 1.9 mmol) are dissolved in ethanol, treated with .5 ml, concentrated HCl and refluxed for 1 hour. The reaction is cooled, quenched with water, pH adjusted to 9 and the product extracted 2x ethyl acetate. The combined organics are washed, dried, filtered, concentrated in vacuo and the
10 crude material chromatographed on 150 mesh SiO₂ with ethyl acetate as the eluent. The free base is dissolved in ether and treated dropwise with concentrated HCl. The HCl salt is collected by filtration to yield 0.15 g
15 (35.6%) of the 7:3 endo/exo isomers as a yellow crystalline solid.

MP = 148-149°C.

¹H NMR (300 MHz, CDCl₃): delta 8.5 (m, 1H), 8.16 (m, 1H), 7.7-7.6 (m, 2H), 7.1-7.0 (d, 1H, J=6Hz), 6.9
20 (m, 1H), 6.75 (m, 1H), 4.7 (m, .7H), 4.35 (m, .3H), 3.85 (bs, 3H), 2.7-1.1 (m, 10H).

HRMS 335.1659 Calcd. for C₂₀H₂₁N₃O₂ 335.1633.

25

30

-88-

EXAMPLE 645-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxy-phenyl]-1,2,3,4-Tetrahydro-2-Pyrimidinone

alpha-aminomethylene-[3-(endo-(bicyclo[2.2.1]-
5 hept-2-yloxy)-4-methoxyphenyl]acetaldehyde of Example 46
(0.60 g, 2.08 mmol) and urea (0.20 g, 3.3 mmol) are
dissolved in 10 ml ethanol, treated with 2 ml of con-
centrated HCl solution and refluxed for 1.5 hours. The
reaction is cooled, neutralized with NH_4OH solution and
10 the product extracted 4x ethyl acetate. The organics
are washed, dried, filtered, concentrated in vacuo and
the resulting residue crystallized from ether to afford
0.38 g (58.6%) of 5-[3-(bicyclo[2.2.1]hept-2-yloxy)-4-
methoxyphenyl]-1,2-dihydro-2-pyrimidinone, identical to
15 the product described in Example 47.

The pyrimidinone (.38 g, 1.22 mmol) is dissolved in
20 ml ethanol and treated with .5 g Raney Nickel, 40 psi
hydrogen and refluxed for 18 hours. The reaction is
cooled, filtered through celite, and the filtrate
20 concentrated in vacuo. The residue is crystallized from
ethyl acetate to afford 0.12 g (31.8%) of the dihydro
pyrimidinone as a crystalline solid comprising a 7:3
mixture of endo/exo norbornyl isomers.

MP = 136-137°C.

25 ^1H NMR (300 MHz, CDCl_3): delta 7.1 (bs, 1H),
6.9-6.7 (m, 3H), 6.4 (d, 1H, $J=5\text{Hz}$), 5.4 (bs, 1H), 4.65
(m, .7H), 4.4 (bs, 2H), 4.25 (m, .3H), 3.85 (s, 3H),
2.7-1.2 (m, 10H).

HRMS 314.1644 (+) Calcd. for $\text{C}_{18}\text{H}_{22}\text{N}_2\text{O}_3$ 314.1630.

30 In like manner the exo isomer of the title compound
is prepared in 60.3% yield from the exo isomer of
5-[3-(exo-bicyclo[2.2.1]hept-2-yloxy)-4-methoxyphenyl]-
1,2-dihydro-2-pyrimidinone (Example 47) by hydrogenation
as described above:

-89-

^1H NMR (300 MHz, $\text{CDCl}_3 + \text{CD}_3\text{OD}$): δ 6.7 (m, 3H), 6.2 (bs, 1H), 4.15 (m, 1H), 3.78 (s, 3H), 3.3 (bs, 2H), 2.4-1.0 (m, 10H).

HRMS 314.1647 (M+) Calcd. for $\text{C}_{18}\text{H}_{22}\text{N}_2\text{O}_3$, 314.1625.

EXAMPLE 65

5-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxy-phenyl]hexahydro-2-Pyrimidinone

The tetrahydro pyrimidinone of Example 64 (.30 g, .882 mmol) is dissolved in 15 ml methanol, treated with .3 g Raney Nickel and hydrogenated at 40 psi for 6 hours. The reaction is filtered through celite dried over MgSO_4 , filtered, concentrated in vacuo and triturated from ether to provide .275 g (98%) of the cyclic urea as a crystalline solid. This material is a 7:3 mixture of endo/exo bicycloalkyl isomers.

MP = $>220^\circ\text{C}$.

^1H NMR (300 MHz, CDCl_3): δ 6.85-6.6 (m, 3H), 5.1 (bs, 2H), 4.65 (m, .7H), 4.25 (m, 3H), 3.9 (bs, 3H), 3.5 (m, 4H), 3.1 (m, 1H), 2.7-1.1 (m, 10H).

MS 316 (M+) Calcd. 316.1787.

By means of this procedure there is prepared from appropriate reactants:

5-[3-(indan-2-yloxy)-4-methoxyphenyl]hexahydro-2-pyrimidinone in 75.6% yield:

MP = $212-214^\circ\text{C}$.

^1H NMR (300 MHz, DMSO): δ 7.2-6.8 (m, 7H), 6.3 (bs, 2H), 5.2 (m, 1H), 3.7 (s, 3H), 3.4-3.2 (m, 7H), 3.0 (m, 2H).

^{13}C NMR (75.43 MHz, DMSO): δ 156.0, 148.4, 146.8, 140.8, 132.9, 126.4, 124.6, 119.5, 113.9, 112.2, 79.2, 77.9, 55.4, 45.5, 36.9.

HRMS 338.1629 (M+) Calcd. for $\text{C}_{20}\text{H}_{22}\text{N}_2\text{O}_3$, 338.1630.

-90-

5-[3-(endo-bicyclo[2.2.1]hept-2-yloxy)-4-methoxy-phenyl]hexahydro-2-pyrimidinone in 66.5% yield:

¹H NMR (300 MHz, CDCl₃): 7.2 (bs, 2H), 6.8-6.7 (m, 3H), 4.6 (m, 1H), 3.8 (s, 3H), 3.4 (m, 4H), 3.15 (m, 1H), 2.6-1.0 (m, 10H).

5

5-[3-(exo-bicyclo[2.2.1]hept-2-yloxy)-4-methoxy-phenyl]hexahydro-2-pyrimidinone in 82% yield:

¹H NMR (300 MHz, CDCl₃): delta 6.8-6.7 (m, 3H), 5.4 (bs, 2H), 4.15 (m, 1H), 3.8 (s, 3H), 3.4 (m, 4H), 3.15 (m, 1H), 2.4-1.0 (m, 10H).

10

HRMS 316.1801 (M+) Calcd. for C₁₈H₂₄N₂O₃, 316.1757.

15

20

25

30

-91-

EXAMPLE 66

4-[3-(Indan-2-yloxy)-4-Methoxyphenyl]-4-Hydroxy-1,2,3,4-Tetrahydro-2-Pyrimidinone and 4-[3-(Indan-2-yloxy)-4-Methoxyphenyl]-1,2-Dihydro-2-Pyrimidinone

5 The indanyl substituted enamino-ketone of Example 59 (1.5 g, 4.45 mmol) and urea (.4 g, 6.68 mmol) are dissolved in 10 ml ethanol and 5 ml 1N HCl and refluxed for 2 hours. The reaction mixture is cooled, quenched with H₂O and neutralized with saturated NaHCO₃ solution. The product is extracted 3x ethyl acetate and the
10 combined organics are washed with brine, dried over MgSO₄, filtered and concentrated in vacuo to provide a crude mixture of two products which are separated by SiO₂ chromatography with ethyl acetate/hexanes as the eluent. Obtained is 0.15 g (9%) of 0.15 g (10%) of
15 crystalline materials.

The 4-hydroxy-1,2,3,4-tetrahydropyrimidinone product. MP = 207-208°C.

¹H NMR (300 MHz, CDCl₃/CD₃OD): delta 7.7-7.5 (m, 3H), 7.25-7.1 (m, 4H), 6.91 (d, 1H, J=6Hz), 5.3 (m, 1H),
20 3.9 (s, 3H), 3.45 (dd, 2H, J=5Hz), 3.3 (dd, 2H, J=12Hz, J=3Hz).

HRMS 352.1454 Calcd. for C₂₀H₂N₂O₄ 352.1423.

The 1,2-dihydropyrimidinone product.

25 MP = > 220°C.

¹H NMR (300 MHz, DMSO): delta 8.1 (m, 1H), 7.85 (m, 2H), 7.4-7.0 (m, 6H), 5.5 (m, 1H), 3.95 (s, 3H), 3.5 (m, 3H), 3.25 (bd, 2H, J=12Hz).

HRMS 334.1332 Calcd. for C₂₀H₁₈N₂O₃ 334.1318.

30

-92-

EXAMPLE 674-[3-(Indan-2-yloxy)-4-Methoxyphenyl]-1,2-Dihydro-2-Pyrimidinone

5 The indanyl substituted enamino-ketone of Example 59 (800 mg, 2.4 mmol) and urea (210 mg, 3.56 mmol) are dissolved in 5 ml of 1N HCl and 15 ml ethanol and refluxed for 4 hours. The reaction is cooled, quenched with H₂O and the product is extracted 3x ethyl acetate. The organics are washed, dried and concentrated. The crude product is crystallized from ethyl acetate to provide 0.32 g (40%) as a beige crystalline solid.

10 This material is identical in all respects to the product described in Example 66.

EXAMPLE 68

15 4-[3-(Indan-2-yloxy)-4-Methoxyphenyl]-hexahydro-2-Pyrimidinone

The pyrimidinone of Example 67 (.32 g, .96 mmol) is dissolved in 15 ml of methanol, treated with Raney Nickel and hydrogenated under 40 psi for 8 hours at room temperature. The mixture is filtered through Celite and the catalyst washed several times with methanol. The organics are concentrated in vacuo and the crude residue triturated from ether to provide 20 mg (58%) of the product as a crystalline solid.

25 MP = 182-183°C.

¹H NMR (300 MHz, CDCl₃): delta 7.4-6.8 (m, 7H), 5.55 (bs, 1H), 5.25 (m, 1H), 5.2 (bs, 1H), 4.6 (m, 1H), 3.9 (s, 3H), 3.6-3.2 (m, 6H), 2.2 (m, 1H), 2.0 (m, 1H).

HRMS 338.1665 Calcd. for C₂₀H₂₂N₂O₃ 338.1631.

30

-93-

EXAMPLE 695-[3-(Indan-2-yloxy)-4-Methoxyphenyl]-1,2,3,4-Tetrahydro-2-Pyrimidinone

5- [3-Indan-2-yloxy)-4-methoxyphenyl]-1,2-dihydro-
2-pyrimidinone (740 mg, 2.2 mmol) is dissolved in 10 ml.
acetic acid cooled in an ice bath and treated with
NaCNBH₃ (140 mg, 2.2 mmol). The reaction is stirred for
2 hours and is worked up by dilution with water and
extraction with ethyl acetate. The collected organics
are washed with water, brine, dried over Na₂SO₄,
filtered and concentrated in vacuo to afford 640 mg
(87.4%) of the title product as a white crystalline
solid.

MP = 198-201°C.

¹H NMR (300 MHz, CDCl₃): delta 7.5 (bs, 1H), 7.2
(m, 4H), 6.85 (m, 3H), 6.38 (bd, 1H), 5.9 (bs, 1H), 4.3
(bs, 2H), 3.8 (s, 3H), 3.4-3.1 (m, 4H).

Similarly, 5-[3-(endo-bicyclo[2.2.1]hept-2-yloxy)-
4-methoxyphenyl]-1,2,3,4-tetrahydro-2-pyrimidinone is
prepared in 12.4% yield from the corresponding 1,2-
dihydropyrimidinone:

MP = 205-208°C.

¹H NMR (300 MHz, DMSO): delta 8.2 (bs, 1H),
6.9-6.5 (m, 5H), 4.6 (m, 1H), 4.2 (bs, 2H), 3.8 (s, 3H),
2.5-1.0 (m, 10H).

-94-

EXAMPLE 70

5-[3-(Exo-Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxy-phenyl]hexahydro-2-Pyrimidinone

(A) 3-[(3-bicyclo[2.2.1]hept-2-yloxy)-4-methoxy-phenyl]glutarnitrile

5

Cyanoacetic acid (18.1 g, 0.213 mmol) and exo-norbornyl isovanillin ether (17.5 g, 71.1 mmol) are dissolved in 80 ml of pyridine and 2 ml of piperidine and are heated to 100°C. for 40 hours. The reaction mixture is cooled to room temperature, poured into 200 ml of water and extracted with 2 x 100 ml ethyl acetate. The combined organics are washed with water, 1N HCl, saturated NaHCO₃ solution, water, dried over MgSO₄, filtered and concentrated in vacuo to provide a brown crystalline residue which is recrystallized from ether to afford 15.1 g (68.5%) of the dicyanide as a crystalline solid.

10

15

MP = 122-123°C.

¹H (300 MHz, CDCl₃): 6.82-6.65 (m, 3H), 4.14 (m, 1H), 3.8 (s, 3H), 3.3 (m, 1H), 2.77 (m, 1H), 2.45 (m, 1H), 2.3 (m, 1H), 1.8-1.0 (m, 8H).

20

25

30

-95-

(B) 3-[(3-bicyclo[2.2.1]hept-2-yloxy)-4-Methoxy-phenyl]glutaramide

The above-prepared glutaronitrile from (A) (14.8 g, 47.7 mmol) is dissolved in 200 ml of acetone and is treated at 0°C. with 100 ml water, 33.8 ml 30% H₂O₂ and 21.2 ml 10% Na₂CO₃. The reaction is warmed slowly and stirred at room temperature for 14 hours. The reaction mixture is concentrated to 150 ml and the residue is partitioned between 100 ml of water and 200 ml of ethyl acetate. The organics are washed with water, dried over MgSO₄, filtered and concentrated in vacuo to provide the crude diamine which is titrated from ether to provide 13.8 g (84%) of the diamide as a crystalline solid.

MP = 175-177°C.

¹H NMR (300 MHz, CDCl₃): 6.8-6.6 (m, 3H), 4.15 (m, 1H), 3.75 (s, 3H), 3.6 (m, 1H), 2.6-1.0 (m, 14H).

(C) 5-[3-(exo-bicyclo[2.2.1]hept-2-yloxy)-4-methoxyphenyl]hexahydro-2-pyrimidinone

The glutaramide (B) (1 g, 2.89 mmol) is dissolved in pyridine. To it is added lead tetraacetate (2.72 g, 6.13 mmol) and the mixture stirred for 20 hours at room temperature. The reaction mixture is diluted with water and extracted 2 x 100 ml ethyl acetate. The organics are washed with brine, dried over MgSO₄, filtered and concentrated in vacuo. The crude product is crystallized from ethyl acetate to provide 0.60 g (65.7%) of the cyclic urea as a crystalline solid.

MP = 191-192°C.

¹H NMR (300 MHz, CDCl₃): 6.8-6.6 (m, 3H), 5.35 (bs, 2H), 4.15 (m, 1H), 3.8 (s, 3H), 3.4 (m, 4H), 3.1 (m, 1H), 2.5-1.0 (m, 10H).

-96-

In like manner analogous formula (I) compounds wherein the bicycloalkyl group (R^1) is as defined below are prepared from appropriately substituted benzaldehydes:

	<u>R^1</u>
5	endo-bicyclo[2.2.1]hept-2-yl
	exo-bicyclo[3.2.1]oct-2-yl
	endo-tricyclo[3.2.1.0 ^{2,6}]dec-4-yl
	exo-tricyclo[3.2.1.0 ^{2,6}]dec-8-yl
10	exo-bicyclo[3.3.1.1 ^{3,7}]dec-2-yl
	indan-2-yl

15

20

25

30

-97-

PREPARATION A3-(Exo-Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxybenzaldehyde

Endo-bicyclo[2.2.1]hept-2-ol (5.6 g, 5.0 mmol), isovanillin (7.6 g, 50.0 mmol) and triphenylphosphine (19.65 g, 75.0 mmol) are dissolved in 250 ml of dry tetrahydrofuran and to this mixture is added dropwise diethylazodicarboxylate (11.80 ml, 75.0 mmol). The reaction mixture is heated to reflux and is allowed to reflux for 48 hours. At this time the reaction mixture is cooled to room temperature and is diluted with 500 ml of diethyl ether. The collected organics are washed with 2 x 200 ml water, 2 x 200 ml .5N sodium hydroxide solution, 1 x 100 ml water and 1 x 100 ml saturated sodium chloride solution. The organics are dried over anhydrous sodium sulfate, filtered and concentrated in vacuo. The crude reaction mixture is chromatographed on SiO₂ (32-60 mesh) with 15% ethyl acetate/hexanes as the eluent. The appropriate fractions are collected and concentrated to yield 5.35 g (43.5%) of the exclusively exo-compound as a clear yellow oil.

¹H NMR (300 MHz, CDCl₃): delta 9.83 (s, 1H), 7.4 (dd, 1H, J=9Hz, J=1Hz), 7.3 (d, 1H, J=1Hz), 6.95 (d, 1H, J=9Hz), 4.28 (m, 1H), 3.91 (s, 3H), 2.6-1.0 (m, 10H).

Similarly, 3-(exo-Tricyclo[3.2.1.0^{2,6}]dec-4-yloxy)-4-methoxybenzaldehyde is prepared from tricyclo[3.2.1.0^{2,6}]decan-4-ol in 68.3% yield.

¹H NMR (300 MHz, CDCl₃): delta 9.8 (m, 1H), 7.4 (m, 2H), 6.95 (m, 1H), 4.95 (m, 1H), 4.0 (s, 3H), 2.8-1.2 (m, 14H).

3-(Endo-tricyclo[3.2.1.0^{2,6}]dec-8-yloxy)-4-methoxybenzaldehyde is prepared from tricyclo[3.2.1.0^{2,6}]decan-8-ol in 60.2% yield.

-98-

¹H NMR (300 MHz, CDCl₃): delta 9.8 (s, 1H), 7.4-7.3 (m, 2H), 6.85 (m, 1H), 4.8 (m, 1H), 3.85 (s, 3H), 2.3-.9 (m, 14H).

PREPARATION B

5 2-(Bicyclo[2.2.1]hept-2-yl)-1,3-Benzodioxole

Norcamphor (25 g, 227 mmol) and pyrocatechol (22.7 g, 206 mmol) are dissolved in 500 ml of toluene and is heated to reflux in the presence of a catalytic amount of para-toluene sulfonic acid. The reaction mixture is refluxed over a soxhlet extractor charged with 3A molecular sieves overnight (15 hours). The reaction is cooled to room temperature and the toluene is removed in vacuo. The residue is dissolved in 500 ml ethyl ether and the ethereal layer is washed with 1 x 100 ml 2N NaOH, solution and 2 x 100 ml water and 1 x 100 ml saturated NaCl solution. The organics are dried over MgSO₄, filtered and concentrated in vacuo to yield 35.2 g (85%) of the catechol ketal as a white solid. MP: 42-43°C.

20 ¹H NMR (300 MHz, CDCl₃): delta 6.8 (m, 4H), 2.5-1.2 (m, 10H).

Following the above procedure tricyclo[3.2.1.0^{2,6}]-decan-8-one is converted to 2-(tricyclo[3.2.1.0^{2,6}]dec-8-yl)-1,3-benzodioxole in 84.8% yield.

25 ¹H NMR (300 MHz, CDCl₃): delta 6.9 (m, 4H), 2.7-1.0 (m, 14H);

benzobicyclo[2.2.1]heptan-2-one is converted to 2-(benzobicyclo[2.2.1]hept-2-yl)-1,3-benzodioxole in 93.5% yield.

30 ¹H NMR (300 MHz, CDCl₃): delta 7.4-7.2 (m, 4H), 6.9-6.7 (m, 4H), 3.6 (m, 2H), 2.6 (m, 1H), 2.4 (m, 1H), 2.25 (m, 1H), 2.05 (m, 1H).

PREPARATION C2-(Endo-Bicyclo[3.2.1]hept-2-yloxy)phenol

Aluminum chloride (4.9 g, 37 mmol) is suspended in 50 ml of dry ether and is cooled to 0°C. To this is added 12.5 ml of a 1M solution of lithium aluminum hydride in ether (12.5 mmol). The slurry is stirred for a half hour and to this is added the 2-(bicyclo[2.2.1]-hept-2-yloxy)-1,3-benzodioxole (5.00 g, 25 mmol) as a solution in 50 ml ethyl ether. The solution is stirred for a half hour and then is carefully quenched dropwise with 50 ml of a saturated solution of sodium potassium tartrate. The resulting slurry is basified to a pH of 12 with 2N NaOH solution and then is back titrated to pH 7 with 10% hydrochloric acid solution. The layers are separated and the aqueous layer was re-extracted 2 x 150 ml ether. The combined organics are washed with 2 x 200 ml water and 1 x 200 ml saturated NaCl solution. The ethereals are dried over magnesium sulfate, filtered and concentrated in vacuo to yield (4.6 g 90%) of the phenol as a clear oil.

¹H NMR (300 MHz, CDCl₃): delta 7.0-6.7 (m, 4H), 6.7 (bs, 1H), 4.7 (m, 1H), 2.68 (m, 1H), 2.36 (m, 1H), 2.14 (m, 1H), 1.96 (m, 1H), 1.66 (m, 1H), 1.46 (m, 4H), 1.22 (m, 1H).

25 2-(Endo-tricyclo[3.2.1.0^{2,6}]dec-8-yloxy)phenol is prepared in 97% yield from 2-(tricyclo[3.2.1.0^{2,6}]dec-8-yl)-1,3-benzodioxole.

¹H NMR (300 MHz, CDCl₃): delta 7.0-6.7 (m, 4H), 5.7 (s, 1H), 4.65 (m, 1H), 2.6-1.0 (m, 14H).

30 2-(Endo-benzobicyclo[2.2.1]hept-2-yloxy)phenol is produced in 92.6% yield from 2-(benzobicyclo[2.2.1]-hept-2-yl)-1,3-benzodioxole:

¹H NMR (300 MHz, CDCl₃): delta 7.35-7.2 (m, 4H), 6.8-7.0 (m, 4H), 5.25 (m, 1H), 4.95 (bs, 1H), 3.77 (m,

-100-

1H), 3.45 (m, 1H), 2.45 (m, 1H), 2.0 (m, 1H), 1.85 (1.2 (m, 1H).

PREPARATION D

4-Bromo-2-(Endo-Bicyclo[2.2.1]hept-2-yloxy)phenol

5 2-(Endo-bicyclo[2.2.1]hept-2-yloxy) (3.93 g, 19.3 mmol) is dissolved in 200 ml of chloroform and is cooled to -20°C. To this is added bromine (3.09 g, 19.3 mmol) as a 1M solution in chloroform dropwise over one half hour. After addition is complete the solution is warmed to room temperature and concentrated in vacuo to yield 10 5.60 g (~100%) of the mono-bromo compound.

¹H NMR (300 MHz, CDCl₃): delta 7.0-6.7 (m, 3H), 5.65 (bs, 1H), 4.55 (m, 1H), 2.6 (m, 1H), 2.3 (m, 1H), 2.1 (m, 1H), 1.85 (m, 1H), 1.6 (m, 1H), 1.4 (m, 4H), 15 1.15 (m, 1H).

 In like manner 2-(endo-tricyclo[3.2.1.0^{2,6}]dec-8-yloxy)phenol is converted to the corresponding 4-bromo derivative in 99% yield.

¹H NMR (300 MHz, CDCl₃): delta 7.0-6.7 (m, 3H), 20 5.75 (s, 1H), 4.65 (m, 1H), 2.7-1.0 (m, 14H).

 2-(endo-benzobicyclo[2.2.1]hept-2-yloxy)phenol to the corresponding 4-bromo phenol in essentially quantitative yield:

¹H NMR (300 MHz, CDCl₃): delta 7.25-6.7 (m, 7H), 25 3.65 (m, 1H), 3.48 (m, 1H), 2.5-1.8 (m, 6H).

-101-

PREPARATION E4-Bromo-2-(Endo-Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxybenzene

5 The endo-bromophenol of Preparation D (5.60 g, 19.9 mmol) is dissolved in 50 ml of dry dimethylformamide and is treated with iodomethane (3.55 g, 25 mmol) and anhydrous potassium carbonate (3.46 g, 25 mmol) and is stirred at room temperature for 15 hours. The solution is poured into 600 ml of a .7 normal NaOH solution. The aqueous layer is extracted 2 x 300 ml ethyl ether and the combined organics are washed 4 x 100 ml water and 1 x 100 ml saturated NaCl solution. The organics are dried over magnesium sulfate, filtered and concentrated in vacuo to yield 5.14 g (87%) of methyl ether.

15 ¹H NMR (300 MHz, CDCl₃): delta 6.9-6.6 (m, 3H), 4.55 (m, 1H), 3.82 (s, 3H), 2.6 (m, 1H), 2.3 (m, 1H), 2.05 (m, 2H), 1.6 (m, 1H), 1.42 (m, 4H), 1.16 (m, 1H).

In like manner, the following compounds are prepared from the appropriate bromophenols:

20 4-Bromo-2-(endo-tricyclo[3.2.1.0^{2,6}]dec-8-yloxy)-4-methoxybenzene:

¹H NMR (300 MHz, CDCl₃): delta 6.9 (m, 2H), 6.6 (m, 1H), 4.5 (m, 1H), 3.8 (s, 3H), 2.7-1.0 (m, 14H).

25 4-bromo-2-(endo-benzobicyclo[2.2.1]hept-2-yloxy)-4-methoxybenzene:

¹H NMR (300 MHz, CDCl₃): delta 7.2-6.9 (m, 6H), 6.6 (m, 1H), 5.0 (m, 1H), 3.65 (m, 1H), 3.50 (s, 3H), 3.35 (m, 1H), 2.4 (m, 1H), 1.85 (m, 1H), 1.75 (m, 1H), 1.2 (m, 1H).

30

-102-

PREPARATION FBicyclo[2.2.2]octan-2-ol

5 Bicyclo[2.2.2]-2-octene (20 g, 185 mmol) is dissolved in 100 ml of dry tetrahydrofuran and cooled to 0°C. To this is added borane-tetrahydrofuran complex (200 mmol) dropwise as a 1M solution in tetrahydrofuran. The reaction is warmed slowly to room temperature where it is stirred for 1 hour. The reaction is quenched slowly with 50 ml of 2N NaOH solution and 100 ml of 30% H₂O₂ and is warmed to 50°C. for 1 hour. The solution is 10 diluted with 500 ml ethyl ether. The aqueous layer is saturated with NaCl and extracted with 100 ml ether. The organics are washed with 1 x 100 ml saturated sodium bisulfite solution, 1 x 100 ml water and 1 x 100 brine. 15 The ethereal layer is dried with anhydrous Na₂SO₄, filtered and concentrated in vacuo to yield 19.27 g (83%) of the bicyclooctanol as a white solid. MP: 210-212°C. ¹H NMR (90 MHz, CDCl₃): delta 3.93 (m, 1H), 2.5 (bs, 1H), 2.3-1.0 (m, 12H). 20

25

30

-103-

PREPARATION GEndo-Tricyclo[3.2.1.0^{2,6}]decan-2-ol

The tricyclodecanone (9.57 g, 63.8 mmol) is dissolved in 350 ml of dry tetrahydrofuran and is cooled to 0°C. The reaction mixture is treated with diisobutyl aluminum hydride as a solution in toluene (48 ml of a 1.5 molar solution, 7 mmol). The reaction is allowed, warmed slowly to room temperature and was stirred at room temperature for 14 hours. The reaction is then diluted with 500 ml of ethyl ether and is quenched with 20 ml of saturated NaK tartrate solution. The layers were separated and the aqueous is re-extracted with 100 ml ethyl ether. The combined organics are washed with 1 x 100 ml pH 7 phosphate buffer, 1 x 100 ml water, 1 x 100 ml brine, dried over MgSO₄, filtered and concentrated in vacuo to yield 9.3 g (95.8%) of the exclusively endo compound as a clear viscous oil.

¹H NMR (300 MHz, CDCl₃): delta 4.15 (m, 1H), 2.45 (m, 1H), 2.2-1.7 (m, 13H).

-104-

PREPARATION H3-(Endo-Tricyclo[3.2.1.0^{2,6}]dec-8-yloxy)
4-Methoxybenzaldehyde

5 The tricyclodecanol of Preparation G (5.17 g, 34.0 mmol), isovanillin (5.17 g, 34.0 mmol) triphenylphosphine (10.7 g, 40.8 mmol) and diethylazodicarboxylate (6.4 ml, 40.8 mmol) are dissolved in 200 ml of tetrahydrofuran and are refluxed together for 36 hours. The reaction is cooled to room temperature and diluted with 400 ml ether and 100 ml water. The layers are separated and dried over MgSO_4 , filtered and concentrated in vacuo. The product is purified by flash chromatography (SiO_2 10% ethyl acetate/hexanes) and the appropriate fractions are concentrated in vacuo to yield 1.28 g (14%) of the tricyclic-aryl aldehyde.

^1H NMR (300 MHz, CDCl_3): delta 9.8 (s, 1H), 7.43 (dd, 1H, $J=9\text{Hz}$, $J=1\text{Hz}$), 7.3 (d, 1H, $J=1\text{Hz}$), 6.9 (d, 1H, $J=9\text{Hz}$), 4.3 (m, 1H), 3.9 (s, 3H), 2.4-.9 (m, 14H).

PREPARATION I

20 3-(Endo-Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxybenzoic Acid

To a solution of 3-(endo-bicyclo[2.2.1]hept-2-yloxy)-4-methoxybenzaldehyde (2.46 g, 10 mmol) in 50 ml of acetone is added Jones reagent (6.5 ml of a 2.5 molar solution 16.9 mmol). The reaction is stirred at room temperature for 2 hours. The reaction mixture is diluted with 100 ml of water and extracted 2 x 50 ml ethyl acetate. The combined organics are washed with brine, water, dried over MgSO_4 , filtered and concentrated in vacuo to yield the crude acid which is crystallized from ether to give 1.9 g (72.5%) of the acid as yellow crystals. MP = 170-171°C.

-105-

PREPARATION JMethyl [3-(Endo-Bicyclo[2.2.1]hept-2-yloxy)-
4-Methoxybenzoate

The 3-(Endo-bicyclo[2.2.1]hept-2-yloxy)-4-methoxy-
benzoic acid of Preparation I (1.9 g, 7.25 mmol) is
dissolved in methanol, treated with concentrated
sulfuric acid (1.1 ml, 10 mmol) and is refluxed for 3
hours. The reaction mixture is cooled to room tempera-
ture, concentrated in vacuo to 15 ml, diluted with 50 ml
ethyl acetate and washed with water, aqueous NaHCO₃,
water dried over MgSO₄ filtered and concentrated to in
vacuo to yield 1.9 g (94.9%) of the ester as a thick
yellow oil.

¹H NMR (60 MHz, CDCl₃): delta 10.0 (bs, 1H),
7.6-7.3 (m, 2H), 6.8-6.6 (m, 1H), 4.6 (m, 1H), 3.8 (s,
3H), 2.7-1.0 (m, 10H).

-106-

PREPARATION K2-(Methylsulfinyl)-1-[3-(Endo-Bicyclo[2.2.1]-
hept-2-yloxy)-4-methoxyphenyl]Ethanone

5 Pentane washed sodium hydride (50% in oil, 72 g, 15 mmol) is suspended in 10 ml of dry dimethylsulfoxide and is heated to 70°C. for 45 minutes. The reaction mixture is cooled to room temperature and diluted with 10 ml tetrahydrofuran. The reaction mixture is cooled to 0°C. and to it is added the ester of Preparation J (1.9 g, 6.9 mmol) dropwise over 20 minutes as a solution in tetrahydrofuran. The reaction mixture is stirred at 5°C. for one half hour and at room temperature for one half hour. The reaction mixture is poured onto 30 ml ice/water, acidified with 1N HCl and the product 10 extracted 3 x 30 ml CH₂Cl₂. The combined organics are dried over MgSO₄, filtered and concentrated in vacuo to yield 1.5 g (67.5%) of the sulfoxide as a viscous yellow oil.

20 ¹H NMR (60 MHz, CDCl₃): delta 7.6-7.2 (m, 2H), 7.0-6.7 (m, 1H), 4.6 (m, 1H), 4.35 (d, J=6Hz), 3.9 (s, 3H), 2.75 (s, 3H), 2.7-1.0 (m, 10H).

25

30

-107-

PREPARATION L2-Hydroxy-2-(Thiomethyl)-1-[3-(Endo-Bicyclo-
[2.2.1]hept-2-yloxy)-4-Methoxyphenyl]Ethanone

5 The sulfoxide of Preparation K (1.5 g, 4.66 mmol)
is dissolved in 15 ml of ethyl acetate and is treated
with 15 ml of 1 normal HCl. The reaction is warmed to
50°C. for 16 hours. The cooled reaction mixture is
quenched with 50 ml of water and extracted 2 x 50 ml
ethyl acetate. The combined organics are washed 4 x
10 water, dried over MgSO_4 , filtered and concentrated in
vacuo to yield 1.48 g (98.6%) of the hydroxy sulfide as
a thick oil.

^1H NMR (60 MHz, CDCl_3): δ 7.7-7.4 (m, 2H),
6.9-6.7 (m, 1H), 4.6 (m, 1H), 3.9 (s, 3H), 2.7-1.0 (m,
15 10H), 2.1 (s, 3H).

PREPARATION M3-(bicyclo[2.2.1]hept-2-yloxy)-4-Methoxyphenylglyoxal

The hydroxy sulfide of Preparation L (1.48 g, 4.6
mmol) is dissolved in 10 ml of chloroform and is treated
20 with $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$. The reaction mixture is stirred for
1.5 hours and is cooled, filtered through celite. The
celite is washed with 3 x chloroform. To the collected
organics is added 20 ml of water and the mixture is
basified with solid Na_2CO_3 . The organics are separated,
25 washed with water, dried over MgSO_4 , filtered and
concentrated in vacuo to yield 1.3 g (96.8%) of the
alpha-keto aldehyde as a thick yellow oil.

^1H NMR (60 MHz, CDCl_3): δ 7.8-7.4 (m, 2H),
7.0-6.7 (m, 1H), 4.8 (m, 2H), 3.9 (s, 3H), 2.8-1.0 (m,
30 10H).

-108-

PREPARATION N3-(Indan-2-yloxy)-4-Methoxybenzaldehyde

1-Indanol (5 g, 37 mmol), isovanillin (7.08 g, 46 mmol) triphenylphosphine (12.21 g, 46 mmol) and
5 diethylazodecarboxylate (8.10 g, 46 mmol) are dissolved
in 250 ml tetrahydrofuran and is refluxed until the
indanol has completely disappeared by thin layer
chromatography. The reaction is cooled and diluted with
100 ml water and is extracted 3 x 50 ml ethyl ether.
10 The collected organics are washed 2 x 50 ml water, 2 x
50 ml 1N NaOH solution, 2 x 50 ml water, 2 x 50 ml pH
7.3 molar phosphate buffer and 2 x 50 ml brine. The
organic layer is dried over MgSO_4 filtered, concentrated
in vacuo and flashed on SiO_2 (32-60 mesh 4:1 1:1
15 hexanes/ethyl acetate) to give 2.29 g (23.1%) of the
indanyl ether as a white solid.

^1H NMR (300 MHz, CDCl_3): delta 9.7 (s, 1H), 7.2
(m, 2H), 7.0-6.9 (m, 4H), 6.73 (m, 1H), 5.05 (m, 1H),
3.7 (s, 1H), 3.25 (dd, 2H, $J=13\text{Hz}$, $J=7\text{Hz}$), 3.15 (dd, 2H,
20 $J=13\text{Hz}$, $J=3\text{Hz}$).

25

30

-109-

PREPARATION O3-(Exo-Benzobicyclo[2.2.1]hept-2-yloxy)-4-methoxybenzaldehyde

The benzonorbornylformate (4 g, .022 mole) is dissolved in 50 ml of toluene and is treated with isovanillin (5.19 g, .034 mole) and a catalytic portion of p-toluenesulfonic acid. The reaction mixture is refluxed for 15 hours, cooled and diluted with 100 ml of ether. The organics are washed with 1N NaOH solution, water, brine, dried over MgSO_4 , filtered and concentrated in vacuo. The crude material is purified by flash chromatography [SiO_2 10 \rightarrow 50% EtOAc/Hexane] to afford .78 g (11.4%) of the aldehyde as a yellow oil.

^1H NMR (300 MHz, CDCl_3): δ 9.7 (s, 1H), 7.4-6.9 (m, 7H), 4.4 (m, 1H), 3.9 (s, 3H), 3.6 (bs, 1H), 3.4 (bs, 1H), 2.25 (m, 1H), 1.95 (m, 3H).

-110-

PREPARATION P3-(Exo-Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxybenzyl Alcohol

5 (3-Exo-bicyclo[2.2.1]hept-2-yloxy)-4-methoxybenz-
aldehyde (3.6 g, 14.6 mmol) is dissolved in 50 ml of
methanol and to it is added portionwise sodium boro-
hydride (.18 g, 4.9 mmol). The reaction is stirred at
25°C. for 45 minutes. The reaction is quenched with 2
ml of water and is concentrated to 25 ml and is
10 partitioned between 50 ml ethyl acetate and 50 ml water.
The organic layer is separated, washed with brine, dried
over MgSO_4 , filtered and concentrated in vacuo to yield
3.37 g (93.1%) of the alcohol as a viscous oil.

^1H NMR (60 MHz, CDCl_3): delta 6.8 (m, 3H), 4.6
15 (bs, 2H), 4.2 (m, 1H), 3.8 (s, 3H), 2.8-1.0 (m, 10H).

Similarly, the following compounds are prepared
from appropriate reactants:

3-(bicyclo[2.2.1]hept-2-yloxy)-4-methoxybenzyl
alcohol is prepared in 94.7% yield as a 7:3 endo/exo
20 mixture of isomers from an isomeric mixture of 3-
(bicyclo[2.2.1]hept-2-yloxy)-4-methoxybenzaldehydes.

^1H NMR (60 MHz, CDCl_3): delta 6.9 (m, 3H), 4.7 (m,
7H), 4.6 (bs, 2H), 4.2 (m, 3H), 3.9 (bs, 3H), 2.6 (bs,
3H), 2.6-1.0 (m, 10H);

25 3-(indan-2-yloxy)-4-methoxybenzyl alcohol in 91%
yield:

^1H NMR (300 MHz, CDCl_3): delta 7.2 (m, 4H),
7.0-6.8 (m, 3H), 5.2 (m, 1H), 4.6 (s, 1H), 3.8 (s, 3H),
3.4 (m, 1H), 3.3 (m, 2H);

30 3-(endo-bicyclo[2.2.1]hept-2-yloxy)-4-methoxybenzyl
alcohol in 98% yield:

^1H NMR (300 MHz, CDCl_3): delta 6.8 (m, 3H), 4.5
(m, 1H), 4.45 (bs, 1H), 3.75 (s, 3H), 2.8 (m, 1H), 2.6
(m, 1H), 2.2-1.0 (m, 8H).

-111-

PREPARATION Q3-(Exo-Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxybenzyl Bromide

5 The alcohol of Preparation P (3.3 g, 13.3 mmol) and carbon tetrabromide (8.83 g, 26.6 mmol) are dissolved in 50 ml of ether and is treated with triphenylphosphine (5.23 g, 19.95 mmol). The reaction is stirred for 2 hours at room temperature. The triphenylphosphine oxide was filtered off and is concentrated in vacuo. The residue is titrated with ethyl acetate. The resulting solid is further purified by flash chromatography with 75% hexanes/ethyl acetate as the eluent. The appropriate fractions are collected and concentrated in vacuo to yield 3.05 g (73.7%) of the bromide as thick yellow oil.

15 ^1H NMR (60 MHz, CDCl_3): delta 6.8 (m, 3H), 5.4 (bs, 2H), 4.2 (m, 1H), 3.8 (s, 3H), 2.6-1.0 (m, 10H).

An approximately 7:3 mixture of endo/exo isomers of 3-(bicyclo[2.2.1]hept-2-yloxy)-4-methoxybenzylbromides is similarly produced from the corresponding carbinols.

20 ^1H NMR (60 MHz, CDCl_3): delta 6.8 (m, 3H), 4.7 (m, 7H), 4.6 (bs, 2H), 4.2 (m, 3H), 3.85 (bs, 3H), 2.6-1.0 (m, 10H).

25

30

-112-

PREPARATION R3-(Exo-Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxybenzyl Cyanide

Sodium cyanide (804 mg, 16.4 mmol) is suspended in 30 ml DMSO and warmed to 70°C. A solution of the bromide of Preparation Q (3 g, 9.65 mmol) in 5 ml DMSO is added to the NaCN/DMSO suspension. The reaction is stirred at this temperature for 1 hour. The reaction mixture is cooled and quenched with 50 ml of water and extracted 3 x 50 ml ether. The combined organics are washed 3 x water, dried over MgSO_4 , filtered and concentrated in vacuo to yield a brown residue which is chromatographed on SiO_2 with 75% hexane/ethyl acetate as the eluent. Concentration in vacuo of the appropriate fractions yielded 1.2 g (48%) of the nitrile as a thick yellow oil.

^1H NMR (60 MHz, CDCl_3): delta 6.9 (m, 3H), 4.25 (m, 1H), 3.85 (s, 2H), 3.7 (s, 2H), 2.65-1.0 (m, 10H).

In like manner an approximately 7:3 endo/exo isomeric mixture of 3-(bicyclo[2.2.1]hept-2-yloxy)-4-methoxybenzyl cyanides is produced from the precursor benzyl bromides.

^1H NMR (300 MHz, CDCl_3): delta 6.9-6.7 (m, 3H), 4.65 (m, 7H), 4.25 (m, 3H), 3.85 (bs, 3H), 3.7 (bs, 2H), 2.65-1.1 (m, 10H).

-113-

PREPARATION S3-(Exo-Tricyclo[3.3.1.1^{3,7}]dec-2-yloxy)-
4-Methoxyphenylbenzaldehyde

Isovanillin (10.75 g, 70.72 mmol), 2-adamantanol
(8.6 g, 56.5 mmol), triphenylphosphine (18.53 g, 70.72
mmol) and diethylazodicarboxylate (11.12 ml, 70.72 mmol)
are dissolved in 280 ml tetrahydrofuran and are warmed
to reflux for 24 hours. The reaction is cooled and
diluted with 500 ml of ether and is washed with 2 x 100
ml water, 2 x 100 ml .5N NaOH solution, 2 x 100 ml
water, 1 x 100 ml 3 molar pH 7 phosphate buffer, 1 x 100
ml brine. The organics are concentrated in vacuo to 100
ml and diluted with 500 ml hexanes to precipitate
triphenylphosphine oxide. The solution is dried over
MgSO₄, filtered and concentrated in vacuo. The pure
aldehyde is isolated by a standard bisulfite
formation/liberation procedure to yield 5.29 g (33%) of
the aldehyde as a viscous oil.

¹H NMR (300 MHz, CDCl₃): delta 7.4 (m, 2H), 6.95
(m, 1H), 4.52 (bs, 1H), 3.92 (s, 3H), 2.3-1.4 (m, 14H).

25

30

-114-

PREPARATION T3-Exo-Tricyclo[3.2.1.0^{2,6}]dec-8-yloxy-4-Methoxybenzaldehyde

Endo-Tricyclo[3.2.1.0^{2,6}]decan-8-ol (5.17 g, 34.0 mmol), isovanillin (5.17 g, 34.0 mmol) triphenylphosphine (10.7 g, 40.8 mmol) and diethylazodicarboxylate (6.4 ml, 40.8 mmol) are dissolved in 200 ml of tetrahydrofuran and are refluxed together for 36 hours. The reaction is cooled to room temperature and diluted with 400 ml ether and 100 ml water. The layers are separated and the organics are washed with 2 x 100 ml H₂O, 2 x .5 normal NaOH, 2 x 100 ml H₂O, 1 x 100 ml brine, dried over MgSO₄, filtered and concentrated in vacuo. The product is purified by flash chromatography (SiO₂ 10% ethyl acetate/hexanes) and the appropriate fractions are concentrated in vacuo to yield 1.28 g (14%) of the tricycli-aryl aldehyde.

¹H NMR (300 MHz, CDCl₃): delta 9.8 (s, 1H), 7.43 (dd, 1H, J=9Hz, J=1Hz), 7.3 (d, 1H, J=1Hz), 6.9 (d, 1H, J=9Hz), 4.3 (m, 1H), 3.9 (s, 3H), 2.4-.9 (m, 14H).

25

30

-115-

PREPARATION U1-[3-(Bicyclo[2.2.1]hept-2-yl)-4-Methoxy-phenyl]ethan-1-ol

The aldehyde (3 g, 12.2 mmol) is dissolved in 10 ml dry THF and is added to a solution of methylmagnesium bromide (15 mmol) in 10 ml of THF. The reaction mixture is allowed to warm slowly to room temperature and is stirred at room temperature overnight. The reaction is quenched with 10 ml aqueous NH_4Cl and the product is extracted 2x ethyl acetate. The combined organics are washed with brine dried over MgSO_4 , filtered and concentrated in vacuo to yield 2.9 g (90.7%) of the carbinol as a 7.3 endo/exo mixture of isomers.

^1H NMR (60 MHz, CDCl_3): delta 7.8-6.8 (m, 3H), 5.0-4.0 (m, 2H), 4.0 (2s, 3H), 3.0-1.2 (m, 13H).

1-[3-(indan-2-yloxy)-4-methoxyphenyl]ethan-1-ol is prepared in like manner from 3-indan-2-yloxy-4-methoxybenzaldehyde in 97.5% yield.

^1H NMR (60 MHz CDCl_3): delta 7.2-6.7 (m, 7H), 5.1 (m, 1H), 4.9 (q, 1H, $J=6\text{Hz}$), 3.8 (s, 3H), 3.3 (m, 4H), 1.5 (d, 3H, $J=6\text{Hz}$).

25

30

-116-

PREPARATION VTricyclo[3.2.1.0^{2,6}]decan-4-ol

Tricyclo[3.2.1.0^{2,6}]decan-4-one (1.5 g, 10 mmol) is dissolved in 25 ml of absolute ethanol and treated with sodium borohydride (0.189 g, 5.0 mmol). The reaction is stirred 24 hours, concentrated in vacuo and the resulting crude is redissolved in ether. The organics are washed 2 x H₂O, 2 x brine, dried, filtered and concentrated in vacuo to afford 1.47 g (96.79%) of the alcohol as a clear oil.

¹H NMR (300 MHz, CDCl₃): delta 4.2 (m, 1H), 2.4-1.2 (m, 14H).

¹³C NMR (75.43 MHz, CDCl₃): delta 76.71, 43.10, 41.75, 39.48, 34.43, 22.32 (6 lines).

-117-

PREPARATION W1-[3-[Bicyclo[2.2.1]hept-2-yloxy]-4-Methoxyphenyl]ethanone

5 The title carbinol of Preparation U (2.9 g, 11.07 mmol) is dissolved in 20 ml of acetone and treated with 12.5 ml of 1 Molar Jones reagent. The reaction is stirred for 2 hours, quenched with water and the product is extracted 3x ethyl acetate. The combined organics are washed with H₂O, brine, dried over MgSO₄, filtered
10 and concentrated in vacuo. The crude product is chromatographed on SiO₂ with 4:1 hexane/ethyl acetate as the eleuent to yield 1.5 g (52.7%) of the methyl ketone as a viscous oil as a mixture of isomers.

¹H NMR (60 MHz, CDCl₃): delta 7.7-7.3 (m, 2H),
15 7.2-6.8 (m, 1H), 5.0-4.1 (m, 1H), 4.0 (bs, 3H), 3.0-1.0 (m, 10H), 2.6 (s, 3H).

1-[3-(Indan-2-yloxy)-4-methoxyphenyl]ethanone is prepared in a similar manner from 1-[3-(indan-2-yloxy)-4-methoxyphenyl]ethan-1-ol in 52.1% yield.

20 ¹H NMR (60 MHz, CDCl₃): delta 7.6-6.7 (m, 7H), 5.2 (m, 1H), 3.9 (s, 3H), 3.3 (m, 4H), 2.4 (s, 3H).

25

30

-118-

PREPARATION X3-(Endo-Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxybenzyl Chloride

5 A solution of 3-(endo-bicyclo[2.2.1]hept-2-yloxy)-4-methoxybenzyl alcohol (8.5 g, 34.3 mmol) and triethylamine (5.7 ml, 41.2 mmol) in methylene chloride (75 ml) is cooled to 0°C. and to it is added methane-sulfonyl chloride (2.9 ml, 37.8 mmol) dropwise. The reaction is then stirred for 2 hours and is worked up by
10 dilution with water and removal of the methylene chloride phase. The aqueous layer is re-extracted with methylene chloride, the combined methylene chloride extracts washed with water and brine, then dried over Na₂SO₄, filtered and concentrated in vacuo to afford 8.0
15 g (87.5%) of the benzyl chloride as an orange oil.

¹H NMR (300 MHz, CDCl₃): delta 6.8 (m, 3H), 4.5 (m, 1H), 4.35 (s, 1H), 3.8 (s, 3H), 2.6-1.0 (m, 10H).

In like manner, 3-(indan-2-yloxy)-4-methoxybenzyl
20 chloride is obtained from the corresponding benzyl alcohol.

HRMS 288.0900 (M+) Calcd. for C₁₇H₁₇O₂Cl, 288.0913.

25

30

-119-

PREPARATION Y3-(Indan-2-yloxy)-4-(Methoxybenzyl Cyanide

3-(Indan-2-yloxy)-4-methoxybenzyl chloride (17.7 g, 61.4 mmol) is dissolved in 150 ml dimethylsulfoxide and is treated with potassium cyanide (7.9 g, 121.5 mmol). The reaction is stirred at room temperature for 4.5 hours then partitioned between water and ethyl acetate. It is washed 2 x H₂O, 2 x brine, dried over MgSO₄, filtered and concentrated in vacuo. The crude material is purified by flash chromatography (SiO₂, 3:1 hexane/ethyl acetate). The appropriate fractions are combined and concentrated in vacuo to afford 9.2 g (53%) of the benzyl cyanide as a pale yellow solid.

¹H NMR (CDCl₃, 300 MHz): delta 7.3-7.2 (m, 4H), 6.9 (m, 3H), 5.22 (m, 1H), 3.9 (s, 1H), 3.75 (s, 2H), 3.5-3.2 (m, 4H).

Similarly, 3-(endo bicyclo[2.2.1]hept-2-yloxy)-4-methoxybenzyl cyanide is prepared in 52% yield from the corresponding benzyl chloride.

¹H NMR (300 MHz, CDCl₃): delta 6.8 (m, 3H), 4.5 (m, 1H), 3.8 (s, 3H), 3.6 (s, 2H), 2.5-1.0 (m, 10H).

PREPARATION Z3-(Bicyclo[2.2.1]hept-2-yloxy)-4-Methoxybenzaldehyde (7:3 Endo:Exo Mixture)

Isovanillin (50 g, .328 mol) is placed in a 1 liter round bottom flask which is equipped with a stir bar and reflux condensor and is charged with 500 ml of dimethylformamide. Potassium carbonate (45.3 g, .328 mol) is added to the reaction mixture which is heated to 80°C. At this temperature exo-2-bromonorbornane (12.57 g, .072 mol, .219 equivalents) is added and the reaction mixture is heated to 120°C. for 48 hours. The reaction is then cooled to room temperature and poured into 300 ml of water. The aqueous layer is extracted with ethyl ether

-120-

(3 x 100 ml). The ethereal layer is washed with 1N sodium hydroxide solution (4 x 100 ml), water (3 x 100 ml) and saturated sodium chloride solution (3 x 100 ml). The organic layer is dried over anhydrous magnesium sulfate, filtered and concentrated in vacuo to yield 15.47 g of a green/brown oil. The crude product is purified by flash chromatography using a gradient eluent system of ethyl acetate/hexane [10% ethyl acetate/hexane → 20% ethyl acetate/hexane] to yield 7.76 g (14.4%) of the aldehyde as a white solid. MP: 75.5°C.-79.5°C.

The material represents a 7:3 mixture of endo:exo bicyclo[2.2.1]hept-2-yl ethers.

¹H-nmr (300 MHz CDCl₃): delta 9.82 (s, 1H), 7.41 (m, 1H), 7.31 (d, 1H), 6.95 (d, 1H, J=9Hz), 4.6-4.7 (m, .7H) [endo], 4.2-4.3 (m, .3H) (exo), 3.93 (s, 2.1H) [endo], 3.91 (s, .9H) [exo], 2.7-.9 (m, 10H).

The following compounds are prepared in like manner from appropriate reactants:

3-(exo-tricyclo[3.2.1.0^{2,6}]dec-4-yloxy)-4-methoxybenzaldehyde in 60.2% yield:

¹H NMR (300 MHz, CDCl₃): delta 9.8 (s, 1H), 7.4-7.3 (m, 2H), 6.85 (m, 1H), 4.8 (m, 1H), 3.85 (s, 3H), 2.3-3.9 (m, 14H).

25

30

-121-

PREPARATION AA3-(Endo-Bicyclo[2.2.1]hept-2-yloxy)-
4-Methoxybenzaldehyde

4-Bromo-2-(endo-bicyclo[2.2.1]hept-2-yloxy)-4-methoxybenzene (5.14 g, 17.4 mmol) is dissolved in 150 ml of dry tetrahydrofuran. This solution is cooled to -78°C. and to it is added dropwise 20.5 ml of a 1.7M solution of tert-butyllithium (34.8 mmol) as a solution in pentane. The solution is stirred at -78°C. for 1.5 hours and is then treated with N,N-dimethylformamide (13.2 ml, 170 mmol) as a solution in 30 ml of tetrahydrofuran. The reaction is stirred at -78°C. for one half hour and is warmed slowly to room temperature. The reaction mixture is diluted with 300 ml ethyl ether and washed with 3 x 200 ml H₂O and 1 x 100 ml brine. The organics are dried over anhydrous magnesium sulfate, filtered and concentrated in vacuo to yield 4.04 g (95%) of the benzaldehyde as a white powder. MP 87-88°C.

¹H NMR (300 MHz, CDCl₃): delta 9.82 (s, 1H), 7.4 (dd, 1H, J=9, J=1), 7.3 (d, 1H, J=1Hz), 6.9 (d, J=9Hz), 4.65 (m, 1H), 3.93 (s, 1H), 2.65 (m, 1H), 2.3 (m, 1H), 2.15 (m, 1H), 2.05 (m, 1H), 1.6 (m, 1H), 1.4 (m, 4H), 1.08 (m, 1H).

Following the above procedure, the following benzaldehyde derivative is produced from the corresponding bromo reactant: 3-(endo-tricyclo-[3.2.1.0^{2,6}]dec-8-yloxy)-4-methoxybenzaldehyde in 64% yield.

¹H NMR (300 MHz, CDCl₃): delta 9.8 (s, 1H), 7.5-7.3 (m, 2H), 6.95 (m, 1H), 4.7 (m, 1H), 3.97 (s, 3H), 2.8-1.0 (m, 14H).

-122-

3-(endo-benzobicyclo[2.2.1]hept-2-yloxy)-4-methoxybenzaldehyde in 51.8% yield:

¹H NMR (300 MHz, CDCl₃): delta 9.6 (s, 1H),
7.3-7.2 (m, 2H), 7.1-6.9 (m, 4H), 6.7 (m, 1H), 5.0 (m,
1H), 3.6 (m, 1H), 3.55 (s, 3H), 3.25 (m, 1H), 2.2 (m,
1H), 1.8 (m, 1H), 1.65 (m, 1H), 1.05 (m, 1H).

PREPARATION BB

3-(Exo-Bicyclo[2.2.2]oct-2-yloxy)-4-Methoxybenzaldehyde

Bicyclo[2.2.2]-2-octanol (5.00 g, 40 mmol),
isovanillin (6.09 g, 40 mmol) and triphenylphosphine
(13.1 g, 50 mmol) are dissolved in dry tetrahydrofuran.
To this reaction mixture is added dropwise diethylazodi-
carboxylate (8.71 g, 50 mmol). The reaction is stirred
for 1 hour at room temperature and is refluxed for 80
hours. The reaction mixture was cooled and concentrated
in vacuo and the residue is mashed 3 x 150 ml of ether
to remove the desired material from the triphenylphos-
phine oxide. The combined ethereal layers are washed
with 2 x 100 ml H₂O, 2 x 100 ml 2N NaOH solution, 2 x
100 ml H₂O and 2 x 100 ml brine. The organics are
dried, filtered and concentrated in vacuo. The compound
is purified by flash silica gel (32-80 mesh) chroma-
tography with 20% ethyl acetate/hexane. The appropriate
fractions are concentrated in vacuo to yield 3.32 g
(32%) of product as a viscous oil.

¹H NMR (250 MHz, CDCl₃): delta 7.4 (dd, 1H,
J=8.5Hz, J=1.5Hz), 7.35 (d, 1H, J=1.5Hz), 6.95 (d, 1H,
J=8.5Hz), 4.52 (m, 1H), 3.91 (s, 3H), 2.2-1.3 (m, 12H).

¹³C NMR (63 MHz, CDCl₃): delta 190.9, 155.7,
147.9, 129.97, 126.2, 112.6, 110.9, 77.5, 56.1, 34.74,
28.18, 25.25, 24.55, 23.29, 22.78, 19.09 (16 lines).

-123-

PREPARATION CC3-(Exo-Bicyclo[3.2.1]oct-2-yloxy)-
4-Methoxybenzaldehyde

5 The cis-bicyclo[3.2.1]-3-octanol (1.90 g, .015 mol)
triphenylphosphine (4.72 g, .018 mol), isovanillin (2.73
g, .018 mol) and diethylazodicarboxylate (2.83 ml, 0.018
mol) are dissolved in 75 ml of dry tetrahydrofuran. The
reaction is refluxed for 15 hours, cooled and diluted
with 50 ml ethyl ether and 50 ml water. The aqueous
10 layer is separated and extracted with 2 x 25 ml ether.
The combined organics are washed with 2 x 25 ml water,
2 x 25 ml 1N NaOH solution, 2 x 25 ml water, 2 x 25 ml
3M pH 7 phosphate buffer and 2 x 25 ml brine. The
organics are dried over MgSO_4 and concentrated in vacuo
15 to 50 ml which is diluted with 250 ml of hexanes and
filtered to remove precipitated triphenylphosphine
oxide. This is repeated 3 times to remove the remainder
of the triphenylphosphine oxide. The resulting organic
layer is dried over MgSO_4 filtered and concentrated in
20 vacuo to yield 1.68 g (43%) of the aryl ether as a clear
oil.

^1H NMR (300 MHz, CDCl_3): δ 10.01 (s, 1H),
7.3-7.1 (m, 2H), 6.8 (m, 1H), 4.4 (m, 1H), 3.78 (s, 3H),
2.3-1.3 (m, 12H).

25

30

-124-

PREPARATION DD3-(Endo-Tricyclo[3.3.1.1^{3,7}]dec-2-yloxy)-
4-Methoxyphenylbenzaldehyde

Isovanillin (10.75 g, 70.72 mmol), 2-adamantanol
5 (8.6 g, 56.5 mmol), triphenylphosphine (18.53 g, 70.72
mmol) and diethylazodicarboxylate (11.12 ml, 70.72 mmol)
are dissolved in 280 ml tetrahydrofuran and are warmed
to reflux for 24 hours. The reaction is cooled and
diluted with 500 ml of ether and is washed with 2 x 100
10 ml water, 2 x 100 ml .5N NaOH solution, 2 x 100 ml
water, 1 x 100 ml 3 molar pH 7 phosphate buffer, 1 x 100
ml brine. The organics are concentrated in vacuo to 100
ml and diluted with 500 ml hexanes to precipitate
triphenylphosphine oxide. The solution is dried over
15 MgSO₄, filtered and concentrated in vacuo. The pure
aldehyde is isolated by a standard bisulfite
formation/liberation procedure to yield 5.29 g (33%) of
the aldehyde as a viscous oil.

¹H NMR (300 MHz, CDCl₃): delta 7.4 (m, 2H), 6.95
20 (m, 1H), 4.52 (bs, 1H), 3.92 (s, 3H), 2.3-1.4 (m, 14H).


25

30

-125-

CLAIMS


1. A compound having the formula




- 5
- 10 wherein R¹ is a polycycloalkyl group having from 7 to 11 carbon atoms;
R² is methyl or ethyl;
X is O or NH; and
Y is a saturated or unsaturated 5- or 6-membered heterocyclyl ring having two nitrogen atoms, said ring
15 being optionally substituted with =O or =S, and wherein when said optional group is present the two nitrogen atoms are in a 1,3-position relative to one another, and said optional group is located on the carbon atom between the two nitrogen atoms; the nitrogen atoms of
20 said heterocyclic rings which bear a hydrogen atom being optionally substituted with (C₁₋₅)alkyl, (C₂₋₅)alkenyl, (C₁₋₅)alkanoyl, benzyl, phenylethyl, benzoyl; 1,2,3-thiadiazolyl 2,2-dioxide, the nitrogen atoms of which are optionally substituted with
25 (C₁₋₅)alkyl, (C₂₋₅)alkenyl, benzyl, phenylethyl, (C₁₋₅)alkanoyl or benzoyl; bicyclic heterocyclyl ring containing a total of three nitrogen atoms, one in each of the rings and one angular nitrogen; a pharmaceutically acceptable acid addition salt of said
30 compound having a basic nitrogen atom; the racemic-diastereomeric mixtures and optical isomers of said compounds.

-126-


2. A compound having the formula




wherein R^1 is a polycycloalkyl group having from 7
to 11 carbon atoms;
 R^2 is methyl or ethyl;
 X O or NH; and
 Y is




(a)




(b)




(c)




(d)




(e)




(f)



(g)









(h)




(i)

-127-



or



R^3 is hydrogen, (C_{1-5}) alkyl, (C_{2-5}) alkenyl, benzyl or phenethyl;

20 R^4 is hydrogen, (C_{1-5}) alkyl, (C_{1-5}) alkanoyl, optionally having a basic group, or benzoyl;

a pharmaceutically acceptable acid addition salt of said compound having a basic nitrogen atom; the racemic-diastereomeric mixtures and optical isomers of said compounds.


25 3. A compound according to claim 2 wherein Y is a 5-membered heterocyclyl ring.

4. A compound according to claim 3 wherein Y is a saturated 5-membered heterocyclyl ring.

30

-128-

5. A compound according to claim 4 wherein Y is



6. A compound according to claim 5 wherein X is

0.

7. A compound according to claim 6 wherein R¹ is a bicycloalkyl group.

8. A compound according to claim 7 wherein R³ is (C₁₋₅)alkyl and R⁴ is hydrogen.

9. A compound according to claim 8 wherein each of R² and R³ is methyl.

10. The compound according to claim 9 wherein R¹ is bicyclo[2.2.1]hept-2-yl.

11. The compound according to claim 9 wherein R¹ is bicyclo[2.2.1]oct-2-yl.

12. The compound according to claim 9 wherein R¹ is bicyclo[3.2.1.]oct-3-yl.

13. The compound according to claim 9 wherein R¹ is tricyclo[3.3.1.1^{5,7}]dec-2-yl.


14. The compound according to claim 9 wherein R¹ is tricyclo[3.2.1.0^{2,6}]dec-8-yl.

15. The compound according to claim 9 wherein R¹ is indan-2-yl.

16. A compound according to claim 2 wherein Y is a 6-membered heterocyclyl ring, said ring being substituted with =oxo.

-129-

17. A compound according to claim 16 wherein Y is



5

18. A compound according to claim 17 wherein X is O.

10 19. A compound according to claim 18 wherein R¹ is bicycloalkyl.


20. A compound according to claim 19 wherein R¹ is bicyclo[2.2.1]hept-2-yl.

15 21. The compound according to claim 20 wherein R² is methyl.

22. A compound according to claim 19 wherein R¹ is indan-2-yl.

23. The compound according to claim 22 wherein R² is methyl.

20 24. A compound according to claim 16 wherein Y is



25

25. A compound according to claim 24 wherein R¹ is bicycloalkyl and X is O.

30 26. A compound according to claim 25 wherein R¹ is bicyclo[2.2.1]hept-2-yl.


27. The compound according to claim 26 wherein R² is methyl.

28. A compound according to claim 19 wherein R¹ is indan-2-yl and X is O.

-130-

29. The compound according to claim 28 wherein R^2 is methyl.

30. A compound according to claim 4 wherein Y is



10 31. A compound according to claim 30 wherein R^1 is bicycloalkyl and X is O.


32. A compound according to claim 31 wherein R^1 is bicyclo[2.2.1]hept-2-yl.

15 33. The compound according to claim 32 wherein each of R^2 and R^3 is methyl; and R^4 is hydrogen.

34. A compound according to claim 31 wherein R^1 is indan-2-yl.

35. The compound according to claim 34 wherein each of R^2 and R^3 is methyl; and R^4 is hydrogen.


20 36. A compound according to claim 2 wherein X is O and Y is



37. A compound according to claim 36 wherein R^1 is bicycloalkyl.


30 38. The compound according to claim 37 wherein R^1 is bicyclo[2.2.1]hept-2-yl; R^2 is methyl; and Y is

-131-




- 5 39. The compound according to claim 37 wherein R^1 is indan-2-yl; and R^2 is methyl; and Y is

10



15

40. A compound according to claim 2 wherein X is O and Y is




20

41. A compound according to claim 40 wherein R^1 is bicycloalkyl.

42. The compound according to claim 41 wherein R^1 is bicyclo[2.2.1]hept-2-yl; R^2 is methyl; and Y is


25



30

43. The compound according to claim 41 wherein R^1 is indan-2-yl; R^2 is methyl; and Y is


-132-



5

44. A compound according to claim 3 wherein Y is

10



45. A compound according to claim 44 wherein Y is bicycloalkyl and X is O.

46. The compound according to claim 45 wherein Y is bicyclo[2.2.1]hept-2-yl and R² is methyl.

47. The compound according to claim 46 wherein Y is indan-2-yl and R² is methyl.


48. A compound according to claim 3 wherein R¹ is tricycloalkyl.

49. A compound according to claim 48 wherein X is O and R² is methyl.

50. A compound according to claim 49 wherein Y is a saturated 5-membered heterocyclic ring.


51. A compound according to claim 50 wherein R¹ is tricyclo[3.3.1.1^{3,7}]dec-2-yl; and Y is

30




52. The compound according to claim 51 wherein Y is

-133-



5 53. A method for treating depression in a human suffering therefrom which comprises administering to said human an anti-depression treating amount of a compound according to claim 2.

10 54. A method according to claim 53 wherein said compound has the formula




15
20 wherein R^1 is a bicycloalkyl group and Y is selected from the group consisting of

25

30


-134-




10



15



20




25

55. A method according to claim 54 wherein R¹ is bicyclo[2.2.1]hept-2-yl or indanyl and R² is methyl.


56. The method according to claim 55 wherein R¹ is bicyclo[2.2.1]hept-2-yl; and Y is

30


-135-




57. The method according to claim 55 wherein R¹ is indan-2-yl and Y is



58. The method according to claim 55 wherein R¹ is bicyclo[2.2.1]hept-2-yl and Y is




59. The method according to claim 55 wherein R¹ is indan-2-yl and Y is



-136-

60. A compound having the formula




10 wherein R¹ is a polycycloalkyl group having from 7 to 10 carbon atoms;

R² is methyl or ethyl; and

R³ is hydrogen, (C₁-₅)alkyl, (C₂-₅)alkenyl, benzyl or phenethyl.

15 61. A compound having the formula



25 wherein R¹ is a polycycloalkyl group having from 7 to 10 carbon atoms;


R² is methyl or ethyl; and

R³ is hydrogen, (C₁-₅)alkyl, (C₂-₅)alkenyl, benzyl or phenethyl.

30


-137-

62. A compound having the formula



10 wherein R^1 is a polycycloalkyl group having from 7 to 10 carbon atoms; and
 R^2 is methyl or ethyl.

63. A compound having the formula




25 wherein R^1 is a polycycloalkyl group having from 7 to 10 carbon atoms; and
 R^2 is methyl or ethyl.

30


-138-

64. A compound having the formula




10 wherein R¹ is a polycycloalkyl group having from 7 to 10 carbon atoms; and
R² is methyl or ethyl.

65. A compound having the formula



20 wherein R¹ is a polycycloalkyl group having from 7 to 10 carbon atoms; and
R² is methyl or ethyl.

66. A compound having the formula




wherein R¹ is a polycycloalkyl group having from 7 to 10 carbon atoms;

-139-

R^2 is methyl or ethyl; and

R^5 is a (C_{1-5}) alkyl group.

67. A compound having the formula




10 wherein R^1 is a polycycloalkyl group having from 7 to 10 carbon atoms;

R^2 is methyl or ethyl; and

R^5 is a $di(C_{1-5})$ alkyl group.

15 68. A compound having the formula




25 wherein R^1 is a polycycloalkyl group having from 7 to 10 carbon atoms; and

R^2 is methyl or ethyl.

30

-140-

69. A compound having the formula



(XVI)

5

wherein R^1 is a polycycloalkyl group having from 7 to
10 10 carbon atoms; and
 R^2 is methyl or ethyl.

70. A pharmaceutical composition for treating
depression in a human suffering therefrom which
comprises an anti-depression treating amount of a
15 compound according to claim 2 and a pharmaceutically
acceptable carrier.

20

25

30

INTERNATIONAL SEARCH REPORT

International Application No PCT/US86/00958

I. CLASSIFICATION F SUBJECT MATTER (If several classification symbols apply, indicate all) ³		
According to International Patent Classification (IPC) or to both National Classification and IPC		
INT. CL ⁴ : C07C 47/11; 69/618; 87/08; 87/29; See Attachment		
U.S. CL : 514/248,274,361,392,399,406,424; See Attachment		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁴		
Classification System	Classification Symbols	
U.S.	514/248,274,361,392,399,406,424 544/281,315,316; 560/81 558/406,408,410 564/342,344,370,372; 568/37,43,425	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁴		
CHEMICAL ABSTRACTS ONLINE STRUCTURE SEARCH		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ¹⁴		
Category ⁵	Citation of Document, ¹⁶ with Indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No. ¹⁸
A	US,A, 3,636,039 (GRUENMAN) 18 January 1972 See entire Document	1-15,48- 52,70
A	US,A, 3,772,288 (HARDTMANN) 13 November 1973 See entire document	1,2,16-29, 53-55,58, 59. 70
A	US,A, 3,923,833 (GRUENMAN) 2 December 1975 See entire document	1-15,48- 52, 70
A	US,A, 4,006,243 (STREHLKE) 1 February 1977 See entire document	1-3
Y	US,A, 4,012,495 (SCHMIECHMEN) 15 March 1977 see entire document	1-4,48- 50,53-55,70
Y	US,A, 4,153,713 (HUTH) 8 May 1979, see entire document	1-4,48-50, 53-55,70
<p>⁵ Special categories of cited documents: ¹⁶</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search ²	Date of Mailing of this International Search Report ²	
July 1, 1986	22 JUL 1986	
International Searching Authority ¹	Signature of Authorized Officer ²⁰	
ISA/US	Stephen M. Kapner	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, ¹⁶ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No ¹⁸
Y	US,A, 4,193,926 (SCHMIECHEN) 18 March 1980 see entire document.	1-4,48- 50,53-55, 70
A	US,A, 4,261,995 (WEINHARDT) 14 April 1981 See entire document	1,2,53,70
Y	US,A, 4,308,278 (SCHNEIDER) 29 December 1981 See entire document	1-15,48- 52, 70
Y	US,A, 4,567,263 (EICKEN) 28 January 1986 see entire document	1,2,40- 43,48,49, 70
Y	US,A, 4,582,834 (STENZEL) 15 April 1986 see entire document	1,2,16- 23,28,29, 70
A	EP,A, 0,134,928 (IRIKURA) 27 March 1985 see entire document	1,2,36- 39,70
Y	DE,A, 3,315,797 (STENZEL) 31 October 1984 see entire document	1,2,16- 23,28,29, 70
A	FR,A, 1,451,154 (LAB. GUY HUE) 25 July 1986 see entire document	1,2,48, 49

ATTACHMENT

I. CLASSIFICATION OF SUBJECT MATTER:

INT. Cl. ⁴	C07C:	121/68, 147/14, 149/273
	C07D	207/263, 231/12, 233/3, 233/42, 233/60, 239/10, 239/22, 239/36, 285/06, 471/02,
	A61K	31/38, 31/40, 31/415, 31/505